

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 February 2005 (10.02.2005)

PCT

(10) International Publication Number
WO 2005/012526 A1

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/47, C12Q 1/68, G01N 33/68

(21) International Application Number:
PCT/CA2004/001449

(22) International Filing Date: 30 July 2004 (30.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/491,968 4 August 2003 (04.08.2003) US

(71) Applicant (for all designated States except US): **THE
HOSPITAL FOR SICK CHILDREN** [CA/CA]; 555
University Avenue, Toronto, Ontario M5G 1X8 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SCHERER,
Stephen, W.** [CA/CA]; 60 Mayfield Avenue, Toronto, Ontario
M6S 1K5 (CA). **MINASSIAN, Berge, A.** [CA/CA];
18 Vanderhoof Avenue, Toronto, Ontario M4G 2H1 (CA).

(74) Agent: **BERESKIN & PARR**; 40 King Street West, Suite
4000, Toronto, Ontario M5H 3Y2 (CA).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SI, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **LAFORA'S DISEASE GENE**

(57) Abstract: A novel gene (EPM2B) that is mutated in humans and dogs with Lafora's disease is described.

WO 2005/012526 A1

B4

- 1 -

TITLE: LAFORA'S DISEASE GENE**FIELD OF THE INVENTION**

The invention relates to a novel gene, *EPM2B*, that is involved in Lafora's disease; the protein, malin, encoded by the gene; and methods of
5 diagnosing and treating Lafora's disease.

BACKGROUND OF THE INVENTION

Lafora's disease (LD, OMIM 254780) is the most common and severe form of adolescent-onset progressive epilepsy. Increasing seizures are paralleled with an insidious cognitive decline towards dementia, and death
10 usually within 10 years of onset (1,2). At the cellular level, LD is characterized by an endoplasmic reticulum (ER)-associated accumulation (3) of starch-like glucose polymers (4) called polyglucosans (or Lafora bodies). Inheritance is autosomal recessive with genetic heterogeneity but the clinical presentation is homogeneous (5). The inventors previously discovered that mutations in the
15 *EPM2A* gene on chromosome 6q24 encoding a dual-specificity phosphatase (named Laforin) with a carbohydrate binding domain, cause LD (6, 7 and WO 00/05405).

There is a need in the art to identify other genes involved in Lafora's disease to assist in the diagnosis and treatment of Lafora's disease.

20 SUMMARY OF THE INVENTION

The present inventors positionally cloned a novel gene, *EPM2B*, on chromosome 6p22.3. It encodes a protein with a putative RING-finger domain and 6 NHL-motifs, which are features of complexes designed for ubiquitin-mediated regulation of specific substrates (8,9) and protein-protein
25 interactions (10-13), respectively. Twenty-one distinct DNA sequence variations in *EPM2B* predicted to cause deleterious effects on the protein product, named malin, were found to co-segregate with LD in 39 families. Both laforin and malin localize to the ER suggesting they operate in a related pathway protecting against neuronal polyglucosan accumulation and epilepsy.
30 The inventors have also isolated and sequenced the canine version of *EPM2B* and have shown a mutation in *EPM2B* in dogs with LD.

-2 -

Accordingly, the present invention provides an isolated nucleic acid molecule that is associated with Lafora's disease and having the sequence shown in SEQ ID NO:1 (Figure 6A) (human *EPM2B*). The present invention also provides an isolated nucleic acid molecule that is associated with

5 Lafora's disease and having a sequence shown in SEQ ID NO:3 (Figure 7A) (canine *EPM2B*).

Preferably, the purified and isolated nucleic acid molecule comprises:

- (a) a nucleic acid sequence as shown in SEQ ID NO:1 (Figure 6A) and SEQ ID NO:3 (Figure 7A), wherein T can also be U;
- 10 (b) a nucleic acid sequence complementary to (a);
- (c) a nucleic acid sequence that has substantial sequence homology to a nucleic acid sequence of (a) or (b);
- (d) a nucleic acid sequence that is an analog of a nucleic acid sequence of (a), (b) or (c); or
- 15 (e) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (a), (b), (c) or (d) under stringent hybridization conditions.

The present invention also includes an isolated protein containing a zinc finger of the RING type and 6 NHL-repeat domains which is associated with Lafora's disease. In a preferred embodiment of the invention, the protein

20 has the amino acid sequence as shown in SEQ ID NO:2 (Figure 6B) (human *EPM2B*). In another embodiment, the protein has the amino acid sequence shown in SEQ ID NO:4 (Figure 7B) (canine *EPM2B*).

As shown in Table 1, the inventors have found 21 different mutations in the human *EPM2B* gene that are associated with Lafora's disease.

25 Accordingly, the present invention provides a method of detecting Lafora's disease comprising detecting a mutation in the *EPM2B* gene in a sample from a mammal, preferably human. In a preferred embodiment, the mutation is one listed in Table 1.

The inventors have also discovered a mutation in the *EPM2B* gene in

30 dogs with Lafora's disease. In particular, all affected dogs studied had a bi-allelic expansion of a dodecamer repeat, termed D, and having the sequence GCCGCCCCCGC that starts at nucleotide number 1001 of canine *EPM2B*

-3 -

sequence shown in SEQ ID NO:3. Accordingly, the invention further provides a method of detecting Lafora's disease in a canid comprising detecting a repeat of the sequence GCCGCCCGCCG (SEQ ID NO:5) which starts at nucleotide number 1001 in the canine sequence of *EPM2B* (SEQ ID NO:3).

- 5 Other features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and
10 scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention will now be described in relation to the drawings in which:

- 15 Figure 1. The *EPM2B* region on 6p22.3. The previous 2.2 Mb critical interval was delimited by microsatellite markers *D6S1567* and *BV012568* (15). The *BV012568* boundary was based on loss of homozygosity in LD individuals from an F-C family (LD6; see Fig. 2). The *D6S1567* telomeric boundary was defined by a recombination occurring between *BV012563* and
20 *D6S1567* in an unaffected sibling of a second consanguineous F-C family (15) (LD27). A break in the chain of homozygosity of markers in the LD38324 family allowed the centromeric boundary to be further re-defined to *D6S1688* (Figure 2) The region contains 7 previously annotated genes and the newly
25 discovered *EPM2B* (Figure 6A), which comprised of a single 1188bp coding exon. Representative human and mouse cDNA sequences are shown, as are the putative ATG start and AATAAA polyadenylation signals. The ATG start follows an in-frame stop (at position -60) and the corresponding AUG is present at the beginning of the predicted ORF. The nucleotide sequence surrounding the start (CGCGCCAUGG) has the proposed features of the
30 consensus sequence (GCCA/GCCAUGG) of an eukaryotic translation initiation site (29). *EPM2B* is predicted to encode a 42.3 kDa (395 aa) protein (malin) containing detectable zinc-binding RING-finger and 6 NHL-repeat

-4 -

domains (FIGURE 6B). The RING and NHL acronyms arise from descriptions of the first proteins identified to carry them, namely, the Really Interesting New Gene 1 in *Homo sapiens* (30) and the NCL-1 (11)/HT2A (10)/LIN-41 (13) genes, respectively. It should be noted that RING- and/or NHL- domains occur in a variety of proteins which can have one or both of cytoplasmic or nuclear localization (8,9,12,18). Malin is the only protein so far described having RING and NHL motifs only (there are other proteins with this combination but they also have other associated motifs such as RING-B-Box-coiled-coil domains). The site of a common C332T (P111L) polymorphism is shown by an asterisk (*) (see Table 1). Malin shares 79%, 80%, and 85% homology with the predicted rat (419 aa), mouse (401 aa) and dog (402 aa) proteins, respectively. The variable amino acids were primarily located in the carboxy- and amino- ends of the protein and not in the RING finger or NHL domains. The microsatellite markers beginning with BV- were generated in this study.

Figure 2. Refinement of the *EPM2B* critical interval by haplotype analysis in LD families and mutations in the *EPM2B* gene. **a**, The centromeric boundary was narrowed to *D6S1688* refining the critical region to 840 kb based on the loss of homozygosity in both probands in family LD38324. **b**, Sequence analysis of *EPM2B* identifies a homozygous 76T>A change in family LD6 (as well as in the LD7, LD27, and LD28 F-C families, Table 1). Affected individuals in family LD38324 were found to be homozygous for a dinucleotide deletion (1048-1049delGA) leading to frame-shift mutation in the fifth NHL domain.

Figure 3. RNA hybridization expression analysis of *EPM2B* in human tissues. **a**, A multiple tissue blot (Clontech) was hybridized with a 557bp fragment of the coding region of *EPM2B*. Two transcripts 2.4 kb and 1.5 kb in size were identified in all tissues. **b**, The same sized transcripts were found in tissues from all regions of the brain tested.

Figure 4. Electron micrograph of brain biopsy material from patient LD32817 (*EPM2B* mutation 98T>C). **A**, axon (note the numerous normal neurotransmitter vesicles); **S**, synapse; **LB**, Lafora body (large rounded

-5 -

structure) composed of a dense accumulation of polyglucosan filaments (PG) completely occupying the dendrite. Bar equals 500nm.

Figure 5. Cellular localization of the malin and laforin proteins. **a**, Myc-tagged malin (construct pcDNA3myc*EPM2B*) forms a distinct reticular pattern
5 around the nucleus, as well as within the nucleus. Co-staining with antibody against the GRP94 endoplasmic reticulum (ER)-specific marker reveals co-localization with malin. **b**, Co-localization of the cytoplasmic isoform of laforin (construct pcDNA3myc*EPM2A* (24)) with the ER-specific marker GRP94.

Figures 6A and B (SEQ ID NOS:1 and 2) provides the human nucleic
10 acid and amino acid sequence of *EPM2B*.

Figures 7A and B (SEQ ID NOS:3 and 4) provides the canine nucleic acid and amino acid sequence for *EPM2B*.

Figure 8 shows the profuse starch-like polyglucosan accumulations (Lafora Bodies; LB) form in tissues of affected MWHD with characteristics and
15 cellular distribution patterns identical to human LD (1,29,30). Affected MWHD tissues included nine muscle and two liver biopsies, as well as skin samples and four whole brains obtained at necropsy. Control tissues did not exhibit LB (seven age-matched animals including two normal MWHD and three non-LD epileptic dogs) and are not shown. Histochemical slides were stained with
20 periodic acid-Schiff following diastase treatment, which allows specific detection of polyglucosans (1). Innumerable LB were present throughout the brain in affected MWHD, located, as in human LD (1), in neurons and not glia and in neuronal perikarya and dendrites and not axons: **a**, LB (arrows) in the molecular layer of cerebral cortex (X100 magnification); **b**, Example of a
25 dendritic LB (asterisk); multiple axons (A) synapsing with this dendrite do not contain LB (bar=500nm). **c**, In liver, LB (arrows) were only in gluconeogenic (periportal) hepatocytes, as in human (29); (PV, portal vein; D, portal ductule; X300). **d**, **e**, In skin, LB (arrows) were in sweat glands, specifically, as in human (30), in myoepithelial cells surrounding apocrine sweat glands (**d**,
30 X100) and duct cells of merocrine sweat glands (**e**, foot pad skin tissue, X250). **f**, In skeletal muscle, LB were within vacuoles surrounded by membranes (arrowheads), again as in human LD (29) (bar=500nm).

-6 -

Figure 9 shows MWHd LD links to the *EPM2B* gene, which contains an expansion mutation in affected dogs. a, Pedigree and haplotypes; numbers in black are genotypes, in descending order, at chromosome CFA35 microsatellites REN256I01, REN282I22, D02001, REN157J09, REN214H22, REN94K23, REN166C14, REN172L08; in red, the position of the *EPM2B* gene and the number of D repeats in its coding polymorphic dodecamer repeat; note: expansion mutations are meiotically unstable (e.g. see transmission of expansion from dog 29 to dog 32); E, expanded alleles with precise repeat number not determined; boxes, genotype homozygosities flanking the mutation. b, Multipoint LOD score analysis.

Figure 10 shows the evolution of the *EPM2B* dodecamer polymorphism with the Canidae family of carnivores, presence of an expansion mutation in a non-MWHd epileptic dog, and the inactivating effect of expansions on *EPM2B* mRNA. a, Canine *EPM2B* dodecamer repeat sequence and orthologues; D, exact repeats; T, imperfect repeat (asterisk); red nucleotides, differences with the canine sequence in the repeat region. b, *EPM2B* expansion mutation associated with myoclonic epilepsy in dogs; W, wild-type; M, affected MWHd; B, affected Basset Hound; C, several unaffected carrier MWHd; arrowheads, normal alleles; all other bands, expanded mutant alleles; note how deamination greatly improves PCR detection of expanded alleles even in the presence of the normal sequence. c, *EPM2B* amounts (normalized against *Gapdh*) in skeletal muscle from affected MWHd and controls using real-time quantitative RT-PCR (SYBR Green detection; Supplementary Methods). d, Evolution of the dodecamer polymorphism; the D sequence is not present in feline species; it is present in all canoids: as a single copy in arctoids, and in polymorphic state with one or two copies in canids; both alleles are shown in heterozygous individuals (two sequence rows); all extant carnivores shown appeared sometime after 10 million years ago (Ma).

DETAILED DESCRIPTION OF THE INVENTION

Previous studies by the inventors suggested that ~70% of LD patients carry recessive mutations in the *EPM2A* gene on chromosome 6q24 (6,7,14). To identify the causative gene(s) in the remaining patients, the inventors

-7 -

initially performed linkage and homozygosity mapping on a subset of the non-*EPM2A* LD families (LD6, LD7, LD27, LD28) originating from a French-Canadian (F-C) isolate (15) (Table 1). This approach led to the localization of a second LD locus (*EPM2B*) to a 2.2 Mb region on chromosome 6p22.3 (Figure 1). All affected individuals from these F-C families were found to be homozygous for a rare haplotype across the entire critical interval (Figure 2a). To further refine the locus, five additional LD families having multiply affected siblings were examined using every microsatellite marker that could be developed from the DNA sequence encompassing the critical region. In four families, all affected individuals were homozygous for all markers across the critical interval. In one family (LD38324), however, the chain of homozygosity in two affected individuals extended only partially into the critical region allowing reduction of the *EPM2B* locus to 840 kb between *D6S1688* and *D6S1567* (Figures 1 and 2a).

15 I. NUCLEIC ACID MOLECULES OF THE INVENTION

As hereinbefore mentioned, the present invention relates to isolated nucleic acid molecules that are involved in Lafora's disease. The term "isolated" refers to a nucleic acid substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors, or other chemicals when chemically synthesized. The term "nucleic acid" is intended to include DNA and RNA and can be either double stranded or single stranded.

Broadly stated, the present invention provides an isolated nucleic acid molecule encoding a protein containing a zinc finger of the RING type in the N-terminal portion and 6 NHL-repeat domains in the C-terminal portion which is associated with Lafora's disease. The isolated nucleic acid molecule is preferably the *EPM2B* gene associated with Lafora's disease. In an embodiment of the invention, the isolated nucleic acid molecule has a sequence as shown in SEQ ID NO:1 (Figure 6A) or SEQ ID NO:3 (Figure 7A).

The nucleic acid sequences shown in SEQ ID NOS:1 and 3 (or Figures 6A and 7A, respectively) as well as the mutated sequences specified in Table 1 or in Example 3 can be collectively referred to herein as "the nucleic acid

-8 -

molecules of the invention". The amino acid sequences shown in SEQ ID NOS:2 and 4 (or Figures 6B and 7B, respectively) as well as the mutated sequences specified in Table 1 can be collectively referred to herein as the "proteins of the invention".

- 5 Preferably, the purified and isolated nucleic acid molecule comprises
- (a) a nucleic acid sequence as shown in SEQ ID NO:1 (Figure 6A) or SEQ ID NO:3 (Figure 7A), wherein T can also be U;
 - (b) a nucleic acid sequence complementary to (a);
 - (c) a nucleic acid sequence that has substantial sequence
 - 10 homology to a nucleic acid sequence of (a) or (b);
 - (d) a nucleic acid sequence that is an analog of a nucleic acid sequence of (a), (b) or (c); or
 - (e) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (a), (b), (c) or (d) under stringent hybridization conditions.
- 15 The term "sequence that has substantial sequence homology" means those nucleic acid sequences which have slight or inconsequential sequence variations from the sequences in (a) or (b), i.e., the sequences function in substantially the same manner and can be used to detect, study or treat Lafora's disease. The variations may be attributable to local mutations or
- 20 structural modifications. Nucleic acid sequences having substantial homology include nucleic acid sequences having at least 65%, more preferably at least 85%, and most preferably 90-95% identity with the nucleic acid sequences as shown in SEQ ID NO:1 or SEQ ID NO:3.
- "Sequence identity" can be calculated according to methods known in
- 25 the art. Sequence identity is most preferably assessed by the algorithm of BLAST version 2.1 advanced search. BLAST is a series of programs that are available online at <http://www.ncbi.nlm.nih.gov/BLAST>. The advanced blast search (<http://www.ncbi.nlm.nih.gov/blast/blast.cgi?Jform=1>) is set to default parameters. (ie Matrix BLOSUM62; Gap existence cost 11; Per residue gap
- 30 cost 1; Lambda ratio 0.85 default). References to BLAST searches are: Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. (1990) "Basic local alignment search tool." J. Mol. Biol. 215:403410; Gish, W. & States, D.J.

-9 -

- (1993) "Identification of protein coding regions by database similarity search." *Nature Genet.* 3:266272; Madden, T.L., Tatusov, R.L. & Zhang, J. (1996) "Applications of network BLAST server" *Meth. Enzymol.* 266:131_141; Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D.J. (1997) "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs." *Nucleic Acids Res.* 25:33893402; Zhang, J. & Madden, T.L. (1997) "PowerBLAST: A new network BLAST application for interactive or automated sequence analysis and annotation." *Genome Res.* 7:649656.
- 10 The term "sequence that hybridizes" means a nucleic acid sequence that can hybridize to a sequence of (a), (b), (c) or (d) under stringent hybridization conditions. Appropriate "stringent hybridization conditions" which promote DNA hybridization are known to those skilled in the art, or may be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y.
- 15 (1989), 6.3.1-6.3.6. The term "stringent hybridization conditions" as used herein means that conditions are selected which promote selective hybridization between two complementary nucleic acid molecules in solution. Hybridization may occur to all or a portion of a nucleic acid sequence molecule. The hybridizing portion is at least 50% the length with respect to
- 20 one of the polynucleotide sequences encoding a polypeptide. In this regard, the stability of a nucleic acid duplex, or hybrids, is determined by the T_m , which in sodium containing buffers is a function of the sodium ion concentration, G/C content of labeled nucleic acid, length of nucleic acid probe (l), and temperature ($T_m = 81.5^\circ\text{C} - 16.6 (\log_{10} [\text{Na}^+]) + 0.41\%(\text{G}+\text{C}) - 600/l$). Accordingly, the parameters in the wash conditions that determine
- 25 hybrid stability are sodium ion concentration and temperature. In order to identify molecules that are similar, but not identical, to a known nucleic acid molecule a 1% mismatch may be assumed to result in about a 1°C decrease in T_m , for example if nucleic acid molecules are sought that have a greater
- 30 than 95% identity, the final wash will be reduced by 5°C . Based on these considerations stringent hybridization conditions can be defined as: hybridization at 5 x sodium chloride/sodium citrate (SSC)/5 x Denhardt's

-10 -

solution/1.0% SDS at T_m (based on the above equation) -5°C , followed by a wash of $0.2 \times \text{SSC}/0.1\%$ SDS at 60°C .

The term "a nucleic acid sequence which is an analog" means a nucleic acid sequence which has been modified as compared to the sequence of (a), (b) or (c) wherein the modification does not alter the utility of the sequence as described herein. The modified sequence or analog may have improved properties over the sequence shown in (a), (b) or (c). One example of a modification to prepare an analog is to replace one of the naturally occurring bases (i.e. adenine, guanine, cytosine or thymidine) of the sequence shown in SEQ ID NO:1 or 3, with a modified base such as such as xanthine, hypoxanthine, 2-aminoadenine, 6-methyl, 2-propyl and other alkyl adenines, 5-halo uracil, 5-halo cytosine, 6-aza uracil, 6-aza cytosine and 6-aza thymine, pseudo uracil, 4-thiouracil, 8-halo adenine, 8-aminoadenine, 8-thiol adenine, 8-thiolalkyl adenines, 8-hydroxyl adenine and other 8-substituted adenines, 8-halo guanines, 8 amino guanine, 8-thiol guanine, 8-thiolalkyl guanines, 8-hydroxyl guanine and other 8-substituted guanines, other aza and deaza uracils, thymidines, cytosines, adenines, or guanines, 5-trifluoromethyl uracil and 5-trifluoro cytosine.

Another example of a modification is to include modified phosphorous or oxygen heteroatoms in the phosphate backbone, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages in the nucleic acid molecule shown in SEQ ID NO:1 or SEQ ID NO:3. For example, the nucleic acid sequences may contain phosphorothioates, phosphotriesters, methyl phosphonates, and phosphorodithioates.

A further example of an analog of a nucleic acid molecule of the invention is a peptide nucleic acid (PNA) wherein the deoxyribose (or ribose) phosphate backbone in the DNA (or RNA), is replaced with a polyamide backbone which is similar to that found in peptides (P.E. Nielsen, et al Science 1991, 254, 1497). PNA analogs have been shown to be resistant to degradation by enzymes and to have extended lives *in vivo* and *in vitro*. PNAs also bind stronger to a complimentary DNA sequence due to the lack of

-11 -

charge repulsion between the PNA strand and the DNA strand. Other nucleic acid analogs may contain nucleotides containing polymer backbones, cyclic backbones, or acyclic backbones. For example, the nucleotides may have morpholino backbone structures (U.S. Pat. No. 5,034,506). The analogs may
5 also contain groups such as reporter groups, a group for improving the pharmacokinetic or pharmacodynamic properties of nucleic acid sequence.

It will be appreciated that the invention includes nucleic acid molecules encoding truncations of proteins of the invention, and analogs and homologs of proteins of the invention and truncations thereof, as described below. It will
10 further be appreciated that variant forms of nucleic acid molecules of the invention which arise by alternative splicing of an mRNA corresponding to a cDNA of the invention are encompassed by the invention.

Isolated and purified nucleic acid molecules having sequences which differ from the nucleic acid sequence of the invention due to degeneracy in
15 the genetic code are also within the scope of the invention. Such nucleic acids encode functionally equivalent proteins but differ in sequence from the above mentioned sequences due to degeneracy in the genetic code.

The invention also includes an isolated nucleic molecule that has a mutation as compared to the nucleic acid molecule shown in SEQ ID NO:1
20 (Figure 6A) or SEQ ID NO:3 (Figure 7A), wherein said mutation is associated with Lafora's disease. In a preferred embodiment, the mutation is selected from one of the mutations shown in Table 1 or the canine mutation described in Example 3.

Nucleic acid molecules from the *EPM2B* gene or mutated forms thereof
25 can be isolated by preparing a labelled nucleic acid probe based on all or part of the nucleic acid sequences as shown in SEQ ID NO:1 (Figure 6A) and SEQ ID NO:3 (Figure 7A) or a mutated sequence shown in Table 1, and using this labelled nucleic acid probe to screen an appropriate DNA library (e.g. a cDNA or genomic DNA library). Nucleic acids isolated by screening of a cDNA or
30 genomic DNA library can be sequenced by standard techniques.

Nucleic acid molecules of the invention can also be isolated by selectively amplifying a nucleic acid using the polymerase chain reaction

-12 -

(PCR) methods and cDNA or genomic DNA. It is possible to design synthetic oligonucleotide primers from the nucleic acid molecules as shown in SEQ ID NO:1 (Figure 6A) and SEQ ID NO:3 (Figure 7A) or a mutated sequence shown in Table 1, for use in PCR. A nucleic acid can be amplified from cDNA
5 or genomic DNA using these oligonucleotide primers and standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. It will be appreciated that cDNA may be prepared from mRNA, by isolating total cellular mRNA by a variety of techniques, for example, by using the guanidinium-thiocyanate extraction procedure of Chirgwin et al., Biochemistry, 18, 5294-5299 (1979). cDNA is then synthesized from the mRNA using reverse transcriptase (for example, Moloney MLV reverse transcriptase available from Gibco/BRL, Bethesda, MD, or AMV reverse transcriptase available from Seikagaku America, Inc., St. Petersburg, FL).

15 An isolated nucleic acid molecule of the invention which is RNA can be isolated by cloning a cDNA encoding a novel protein of the invention into an appropriate vector which allows for transcription of the cDNA to produce an RNA molecule which encodes the malin protein. For example, a cDNA can be cloned downstream of a bacteriophage promoter, (e.g. a T7 promoter) in a
20 vector, cDNA can be transcribed *in vitro* with T7 polymerase, and the resultant RNA can be isolated by standard techniques.

A nucleic acid molecule of the invention may also be chemically synthesized using standard techniques. Various methods of chemically synthesizing polydeoxynucleotides are known, including solid-phase synthesis
25 which, like peptide synthesis, has been fully automated in commercially available DNA synthesizers (See e.g., Itakura et al. U.S. Patent No. 4,598,049; Caruthers et al. U.S. Patent No. 4,458,066; and Itakura U.S. Patent Nos. 4,401,796 and 4,373,071).

The initiation codon and untranslated sequences of the nucleic acid
30 molecules of the invention may be determined using currently available computer software designed for the purpose, such as PC/Gene (IntelliGenetics Inc., Calif.). Regulatory elements can be identified using

-13 -

conventional techniques. The function of the elements can be confirmed by using these elements to express a reporter gene which is operatively linked to the elements. These constructs may be introduced into cultured cells using standard procedures. In addition to identifying regulatory elements in DNA, 5 such constructs may also be used to identify proteins interacting with the elements, using techniques known in the art.

The sequence of a nucleic acid molecule of the invention may be inverted relative to its normal presentation for transcription to produce an antisense nucleic acid molecule. Preferably, an antisense sequence is 10 constructed by inverting a region preceding the initiation codon or an unconserved region. In particular, the nucleic acid sequences contained in the nucleic acid molecules of the invention or a fragment thereof, preferably a nucleic acid sequence shown in SEQ ID NO:1 (Figure 6A) and SEQ ID NO:3 (Figure 7A) or a mutated sequence shown in Table 1 may be inverted relative 15 to its normal presentation for transcription to produce antisense nucleic acid molecules.

The antisense nucleic acid molecules of the invention or a fragment thereof, may be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability 20 of the molecules or to increase the physical stability of the duplex formed with mRNA or the native gene e.g. phosphorothioate derivatives and acridine substituted nucleotides. The antisense sequences may be produced biologically using an expression vector introduced into cells in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense 25 sequences are produced under the control of a high efficiency regulatory region, the activity of which may be determined by the cell type into which the vector is introduced.

The invention also provides nucleic acids encoding fusion proteins comprising a novel protein of the invention and a selected protein, or a 30 selectable marker protein (see below).

II. NOVEL PROTEINS OF THE INVENTION

-14 -

The invention further includes an isolated protein encoded by the nucleic acid molecules of the invention. Within the context of the present invention, a protein of the invention may include various structural forms of the primary protein which retain biological activity.

- 5 Broadly stated, the present invention provides an isolated protein containing a zinc finger of the RING type in the N-terminal half and 6 NHL-repeat domains in the C-terminal direction which is associated with Lafora's disease. Preferably, the zinc-binding RING-finger motif (C-X₂-C-X₁₆-C-X₁-H-X₂-C-X₂-C-X₁₄-C-X₂-C) is located between residues 26-71 of the malin protein
10 shown in Figure 6B. The presence of a RING finger is predictive of an E3 ubiquitin ligase function. Therefore, in a preferred embodiment, the protein has a ubiquitin ligase function.

- In a specific embodiment of the invention, the protein has the amino acid sequence as shown in SEQ ID NO:2 (Figure 6B) (human *EPM2B*). In
15 another embodiment, the protein has the amino acid sequence shown in SEQ ID NO:4 (Figure 7B) (canine *EPM2B*).

- In addition to full length amino acid sequences the proteins of the present invention also include truncations of the protein, and analogs, and homologs of the protein and truncations thereof as described herein.
20 Truncated proteins may comprise peptides of at least fifteen amino acid residues.

- Analogous of the protein having the amino acid sequence shown in SEQ ID NO:2 (Figure 6B) or SEQ ID NO:4 (Figure 7B) and/or truncations thereof as described herein, may include, but are not limited to an amino acid
25 sequence containing one or more amino acid substitutions, insertions, and/or deletions. Amino acid substitutions may be of a conserved or non-conserved nature. Conserved amino acid substitutions involve replacing one or more amino acids of the proteins of the invention with amino acids of similar charge, size, and/or hydrophobicity characteristics. When only conserved
30 substitutions are made the resulting analog should be functionally equivalent. Non-conserved substitutions involve replacing one or more amino acids of the

-15 -

amino acid sequence with one or more amino acids which possess dissimilar charge, size, and/or hydrophobicity characteristics.

One or more amino acid insertions may be introduced into the amino acid sequences shown in SEQ ID NO:2 (Figure 6B) or SEQ ID NO:4 (Figure 5 7B). Amino acid insertions may consist of single amino acid residues or sequential amino acids ranging from 2 to 15 amino acids in length. For example, amino acid insertions may be used to destroy target sequences so that the protein is no longer active. This procedure may be used *in vivo* to inhibit the activity of a protein of the invention.

10 Deletions may consist of the removal of one or more amino acids, or discrete portions from the amino acid sequence shown in SEQ ID NO:2 (Figure 6B) or SEQ ID NO:4 (Figure 7B). The deleted amino acids may or may not be contiguous. The lower limit length of the resulting analog with a deletion mutation is about 10 amino acids, preferably 100 amino acids.

15 Analogs of a protein of the invention may be prepared by introducing mutations in the nucleotide sequence encoding the protein. Mutations in nucleotide sequences constructed for expression of analogs of a protein of the invention must preserve the reading frame of the coding sequences. Furthermore, the mutations will preferably not create complementary regions 20 that could hybridize to produce secondary mRNA structures, such as loops or hairpins, which could adversely affect translation of the receptor mRNA.

Mutations may be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the 25 resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site specific mutagenesis procedures may be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. 30 Deletion or truncation of a protein of the invention may also be constructed by utilizing convenient restriction endonuclease sites adjacent to the desired deletion. Subsequent to restriction, overhangs may be filled in, and the DNA

-16 -

religated. Exemplary methods of making the alterations set forth above are disclosed by Sambrook et al (Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, 1989).

The proteins of the invention also include homologs of the amino acid
5 sequence shown in SEQ ID NO:2 (Figure 6B) or SEQ ID NO:4 (Figure 7B)
and/or truncations thereof as described herein. Such homologs are proteins
whose amino acid sequences are comprised of amino acid sequences that
hybridize under stringent hybridization conditions (see discussion of stringent
10 hybridization conditions herein) with a probe used to obtain a protein of the
invention. Preferably, homologs of a protein of the invention will have a
tyrosine phosphatase region which is characteristic of the protein.

A homologous protein includes a protein with an amino acid sequence
having at least 70%, preferably 80-90% identity with the amino acid sequence
as shown in SEQ ID NO:2 (Figure 6B) or SEQ ID NO:4 (Figure 7B).
15 Sequence identity is as previously defined herein.

The invention also contemplates isoforms of the proteins of the
invention. An isoform contains the same number and kinds of amino acids as
a protein of the invention, but the isoform has a different molecular structure.
The isoforms contemplated by the present invention are those having the
20 same properties as a protein of the invention as described herein.

The invention also includes an isolated protein that has a mutation as
compared to the amino acid sequence shown in SEQ ID NO:2 (Figure 6B) or
SEQ ID NO:4 (Figure 7B), wherein said mutation is associated with Lafora's
disease. In a preferred embodiment, the mutation is selected from one of the
25 mutations shown in Table 1 or the canine mutation described in Example 3.

The present invention also includes a protein of the invention
conjugated with a selected protein, or a selectable marker protein (see below)
to produce fusion proteins. Additionally, immunogenic portions of a protein of
the invention are within the scope of the invention.

30 The proteins of the invention (including truncations, analogs, mutants
etc.) may be prepared using recombinant DNA methods. Accordingly, the
nucleic acid molecules of the present invention having a sequence which

-17 -

encodes a protein of the invention may be incorporated in a known manner into an appropriate expression vector which ensures good expression of the protein. Possible expression vectors include but are not limited to cosmids, plasmids, or modified viruses (e.g. replication defective retroviruses, adenoviruses and adeno-associated viruses), so long as the vector is compatible with the host cell used. The expression vectors are "suitable for transformation of a host cell", means that the expression vectors contain a nucleic acid molecule of the invention and regulatory sequences selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid molecule. Operatively linked is intended to mean that the nucleic acid is linked to regulatory sequences in a manner which allows expression of the nucleic acid.

The invention therefore contemplates a recombinant expression vector of the invention containing a nucleic acid molecule of the invention, or a fragment thereof, and the necessary regulatory sequences for the transcription and translation of the inserted protein-sequence. Suitable regulatory sequences may be derived from a variety of sources, including bacterial, fungal, or viral genes (For example, see the regulatory sequences described in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990). Selection of appropriate regulatory sequences is dependent on the host cell chosen, and may be readily accomplished by one of ordinary skill in the art. Examples of such regulatory sequences include: a transcriptional promoter and enhancer or RNA polymerase binding sequence, a ribosomal binding sequence, including a translation initiation signal. Additionally, depending on the host cell chosen and the vector employed, other sequences, such as an origin of replication, additional DNA restriction sites, enhancers, and sequences conferring inducibility of transcription may be incorporated into the expression vector. It will also be appreciated that the necessary regulatory sequences may be supplied by the native protein and/or its flanking regions.

The invention further provides a recombinant expression vector comprising a DNA nucleic acid molecule of the invention cloned into the

-18 -

expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression, by transcription of the DNA molecule, of an RNA molecule which is antisense to a nucleotide sequence comprising the nucleotides as shown

5 SEQ ID NO:1 or SEQ ID NO:3. Regulatory sequences operatively linked to the antisense nucleic acid can be chosen which direct the continuous expression of the antisense RNA molecule.

The recombinant expression vectors of the invention may also contain a selectable marker gene which facilitates the selection of host cells

10 transformed or transfected with a recombinant molecule of the invention. Examples of selectable marker genes are genes encoding a protein such as G418 and hygromycin which confer resistance to certain drugs, β -galactosidase, chloramphenicol acetyltransferase, or firefly luciferase. Transcription of the selectable marker gene is monitored by changes in the

15 concentration of the selectable marker protein such as β -galactosidase, chloramphenicol acetyltransferase, or firefly luciferase. If the selectable marker gene encodes a protein conferring antibiotic resistance such as neomycin resistance transformant cells can be selected with G418. Cells that have incorporated the selectable marker gene will survive, while the other

20 cells die. This makes it possible to visualize and assay for expression of recombinant expression vectors of the invention and in particular to determine the effect of a mutation on expression and phenotype. It will be appreciated that selectable markers can be introduced on a separate vector from the nucleic acid of interest.

25 The recombinant expression vectors may also contain genes which encode a fusion moiety which provides increased expression of the recombinant protein; increased solubility of the recombinant protein; and aid in the purification of a target recombinant protein by acting as a ligand in affinity purification. For example, a proteolytic cleavage site may be added to

30 the target recombinant protein to allow separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein.

-19 -

Recombinant expression vectors can be introduced into host cells to produce a transformed host cell. The term "transformed host cell" is intended to include prokaryotic and eukaryotic cells which have been transformed or transfected with a recombinant expression vector of the invention. The terms "transformed with", "transfected with", "transformation" and "transfection" are intended to encompass introduction of nucleic acid (e.g. a vector) into a cell by one of many possible techniques known in the art. Prokaryotic cells can be transformed with nucleic acid by, for example, electroporation or calcium-chloride mediated transformation. Nucleic acid can be introduced into mammalian cells via conventional techniques such as calcium phosphate or calcium chloride co precipitation, DEAE-dextran-mediated transfection, lipofectin, electroporation or microinjection. Suitable methods for transforming and transfecting host cells can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory textbooks.

Suitable host cells include a wide variety of prokaryotic and eukaryotic host cells. For example, the proteins of the invention may be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus), yeast cells or mammalian cells. Other suitable host cells can be found in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1991).

The proteins of the invention may also be prepared by chemical synthesis using techniques well known in the chemistry of proteins such as solid phase synthesis (Merrifield, 1964, J. Am. Chem. Assoc. 85:2149-2154) or synthesis in homogenous solution (Houbenweyl, 1987, Methods of Organic Chemistry, ed. E. Wansch, Vol. 15 I and II, Thieme, Stuttgart).

III. APPLICATIONS

The present invention includes all uses of the nucleic acid molecule and proteins of the invention including, but not limited to, the preparation of antibodies and antisense oligonucleotides, the preparation of experimental systems to study *EPM2B* and mutated forms thereof, the isolation of substances that modulate *EPM2B* expression and/or activity as well as the

-20 -

use of the *EPM2B* nucleic acid sequences and proteins and modulators thereof in diagnostic and therapeutic applications. Some of the uses are further described below.

A. Diagnostic Applications

5 As previously mentioned, the present inventors have determined that the gene *EPM2B* is mutated in people and canids with Lafora's disease. As a result, the present invention also includes a method of detecting Lafora's disease by detecting a mutation in the *EPM2B* gene or protein.

10 The term "mutation" means any change or difference in the nucleic acid or protein sequence of *EPM2B* as compared to the wild type sequence. Mutations include, but are not limited to, nonsense mutations, missense mutations, frameshift mutations, rearrangement mutations, insertion mutations and deletion mutations.

The sample can be any sample containing the *EPM2B* gene or protein
15 including, but not limited to, biological fluids (such as blood, urine, cerebrospinal fluid, tears, saliva), tissues, tissue extracts, cells and cell extracts.

(i) Detecting Mutations in the Nucleic Acid Sequence

In one embodiment, the present invention provides a method for
20 detecting Lafora's disease comprising detecting a mutation in the *EPM2B* gene in a sample obtained from an animal, preferably a mammal, more preferably a human or canid.

Table 1 summarizes some of the mutations found in human *EPM2B* in patients with Lafora's disease. To date, 21 different DNA sequence
25 alterations have been found in *EPM2B* in 39 families including 8 deletions and 1 Insertion leading to frame-shifts, 8 missense, and 1 non-sense change. Screening assays can be developed for each of the mutations.

The most common mutation identified in seven families is a homozygous 205C→G transition resulting in a proline to alanine change in the
30 RING-finger domain. Accordingly, in one embodiment, the present invention provides a method of detecting Lafora's disease comprising detecting a C → G mutation at position 205 in the *EPM2B* gene (SEQ ID NO:1).

-21 -

Another mutation observed in the four consanguineous F-C families used in the original linkage study all carried a homozygous 76T→A change producing a cysteine-to-serine alteration in one of the 7 conserved cysteine residues that are critical for the zinc-binding ability of the RING-finger domain.

- 5 Accordingly, in another embodiment, the present invention provides a method of detecting Lafora's disease comprising detecting a T → A mutation at position 76 in the *EPM2B* gene (SEQ ID NO:1).

- Another mutation was observed in the LD38324 family that was critical in refining the *EPM2B* locus (Figure 2a) which was a homozygous 2-bp
10 deletion (1048-1049delGA) leading to a frame-shift mutation in the fifth NHL-domain (Figure 2b). Accordingly, in a further embodiment, the present invention provides a method of detecting Lafora's disease comprising detecting a deletion of GA at positions 1048 and 1049 in the *EPM2B* gene (SEQ ID NO:1).

- 15 As described in Example 3, the inventors have also discovered a mutation in the *EPM2B* gene in dogs with Lafora's disease. In particular, all affected dogs studied had a bi-allelic expansion of a dodecamer repeat, termed D, and having the sequence GCCGCCCCCCGC that starts at nucleotide number 1001 of canine *EPM2B* sequence shown in SEQ ID NO:3.
- 20 The inventors have shown that this 12 nucleotide repeat is specific to the canid superfamily, which includes dogs, wolves, foxes, coyotes, and jackals, and have shown that this repeat predisposes dogs to a massive sequence expansion, which is destructive to the *EPM2B* gene and causes Lafora disease. The inventors have thus discovered that canids, including dogs, are
- 25 predisposed to Lafora Epilepsy. Accordingly, the invention further provides a method of detecting Lafora's disease in a canid comprising detecting a repeat of the sequence GCCGCCCCCCGC (SEQ ID NO:5) which starts at nucleotide number 1001 in the canine sequence of *EPM2B* (SEQ ID NO:3). In one embodiment, the method involves detecting at least 3 repeats,
- 30 preferably at least 10 repeats, more preferably from about 14 to about 26 repeats in SEQ ID NO:5.

-22 -

One skilled in the art will appreciate that many suitable methods, in addition to and including the ones discussed in the examples, can be used to detect mutations in the *EPM2B* gene. Detection methods that can be used include, but are not limited to, polymerase chain reaction (PCR), reverse transcriptase PCR, direct sequencing electrophoretic mobility, nucleic acid hybridization, fluorescent in situ hybridization, denaturing high performance liquid chromatography, DNA chip technologies and mass spectroscopy. In one example, in order to isolate nucleic acids from the Lafora's disease gene in a sample, one can prepare nucleotide probes from the nucleic acid sequences of the invention. In addition, the nucleic acid probes described herein can also be used. A nucleotide probe may be labelled with a detectable marker such as a radioactive label which provides for an adequate signal and has sufficient half life such as ^{32}P , ^3H , ^{14}C or the like. Other detectable markers which may be used include antigens that are recognized by a specific labelled antibody, fluorescent compounds, enzymes, antibodies specific for a labelled antigen, and chemiluminescent compounds. An appropriate label may be selected having regard to the rate of hybridization and binding of the probe to the nucleotide to be detected and the amount of nucleotide available for hybridization.

Accordingly, the present invention also relates to a method of detecting the presence of a nucleic acid molecule from the *EPM2B* gene in a sample comprising contacting the sample under hybridization conditions with one or more of nucleotide probes which hybridize to the nucleic acid molecules and are labelled with a detectable marker, and determining the degree of hybridization between the nucleic acid molecule in the sample and the nucleotide probes.

Hybridization conditions which may be used in the methods of the invention are known in the art and are described for example in Sambrook J, Fritsch EF, Maniatis T. In: Molecular Cloning, A Laboratory Manual, 1989. (Nolan C, Ed.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. The hybridization product may be assayed using techniques known in the art. The nucleotide probe may be labelled with a detectable marker as

-23 -

described herein and the hybridization product may be assayed by detecting the detectable marker or the detectable change produced by the detectable marker.

Prior to hybridizing a sample with DNA probes, the sample can be
5 treated with primers that flank the *EPM2B* gene in order to amplify the nucleic acid sequences in the sample. The primers used may be the ones described in the present application. In addition, the sequence of the *EPM2B* gene provided herein also permits the identification and isolation, or synthesis of
10 new nucleotide sequences which may be used as primers to amplify a nucleic acid molecule of the invention, for example in the polymerase chain reaction (PCR) which is discussed in more detail below. The primers may be used to amplify the genomic DNA of other species. The PCR amplified sequences can be examined to determine the relationship between the genes of various species.

15 The length and bases of the primers for use in the PCR are selected so that they will hybridize to different strands of the desired sequence and at relative positions along the sequence such that an extension product synthesized from one primer when it is separated from its template can serve as a template for extension of the other primer into a nucleic acid of defined
20 length. Primers which may be used in the invention are oligonucleotides i.e. molecules containing two or more deoxyribonucleotides of the nucleic acid molecule of the invention which occur naturally as in a purified restriction endonuclease digest or are produced synthetically using techniques known in the art such as for example phosphotriester and phosphodiester methods
25 (See Good et al Nucl. Acid Res 4:2157, 1977) or automated techniques (See for example, Conolly, B .A. Nucleic Acids Res. 15:15(7): 3131, 1987). The primers are capable of acting as a point of initiation of synthesis when placed under conditions which permit the synthesis of a primer extension product which is complementary to the DNA sequence of the invention i.e. in the
30 presence of nucleotide substrates, an agent for polymerization such as DNA polymerase and at suitable temperature and pH. Preferably, the primers are sequences that do not form secondary structures by base pairing with other

-24 -

copies of the primer or sequences that form a hair pin configuration. The primer preferably contains between about 7 and 25 nucleotides.

The primers may be labelled with detectable markers which allow for detection of the amplified products. Suitable detectable markers are
5 radioactive markers such as P-32, S-35, I-125, and H-3, luminescent markers such as chemiluminescent markers, preferably luminol, and fluorescent markers, preferably dansyl chloride, fluorcein-5-isothiocyanate, and 4-fluor-7-nitrobenz-2-axa-1,3 diazole, enzyme markers such as horseradish peroxidase, alkaline phosphatase, β -galactosidase, acetylcholinesterase, or
10 biotin.

It will be appreciated that the primers may contain non-complementary sequences provided that a sufficient amount of the primer contains a sequence which is complementary to a nucleic acid molecule of the invention or oligonucleotide fragment thereof, which is to be amplified. Restriction site
15 linkers may also be incorporated into the primers allowing for digestion of the amplified products with the appropriate restriction enzymes facilitating cloning and sequencing of the amplified product.

In an embodiment of the invention a method of determining the presence of a nucleic acid molecule of the invention is provided comprising
20 treating the sample with primers which are capable of amplifying the nucleic acid molecule or a predetermined oligonucleotide fragment thereof in a polymerase chain reaction to form amplified sequences, under conditions which permit the formation of amplified sequences and, assaying for amplified sequences.

The polymerase chain reaction refers to a process for amplifying a target nucleic acid sequence as generally described in Innis et al, Academic Press, 1990 in Mullis et al., U.S. Pat. No. 4,863,195 and Mullis, U.S. Patent No. 4,683,202 which are incorporated herein by reference. Conditions for amplifying a nucleic acid template are described in M.A. Innis and D.H.
30 Gelfand, PCR Protocols, A Guide to Methods and Applications M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White eds, pp3-12, Academic Press 1989, which is also incorporated herein by reference.

-25 -

The amplified products can be isolated and distinguished based on their respective sizes using techniques known in the art. For example, after amplification, the DNA sample can be separated on an agarose gel and visualized, after staining with ethidium bromide, under ultra violet (UV) light.

- 5 DNA may be amplified to a desired level and a further extension reaction may be performed to incorporate nucleotide derivatives having detectable markers such as radioactive labelled or biotin labelled nucleoside triphosphates. The primers may also be labelled with detectable markers as discussed above. The detectable markers may be analyzed by restriction and electrophoretic
10 separation or other techniques known in the art.

- The conditions which may be employed in the methods of the invention using PCR are those which permit hybridization and amplification reactions to proceed in the presence of DNA in a sample and appropriate complementary hybridization primers. Conditions suitable for the polymerase chain reaction
15 are generally known in the art. For example, see M.A. Innis and D.H. Gelfand, PCR Protocols, A guide to Methods and Applications M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White eds, pp3-12, Academic Press 1989, which is incorporated herein by reference. Preferably, the PCR utilizes polymerase obtained from the thermophilic bacterium *Thermus aquaticus* (Taq
20 polymerase, GeneAmp Kit, Perkin Elmer Cetus) or other thermostable polymerase may be used to amplify DNA template strands.

- It will be appreciated that other techniques such as the Ligase Chain Reaction (LCR) and NASBA may be used to amplify a nucleic acid molecule of the invention (Barney in "PCR Methods and Applications", August 1991,
25 Vol.1(1), page 5, and European Published Application No. 0320308, published June 14, 1989, and U.S. Serial NO. 5,130,238 to Malek).

- In some embodiments of the present invention, mutations in the sequences of the invention may be detected by the direct sequencing of nucleic acid molecules. Techniques for the direct sequencing of DNA are well
30 known in the art. In one embodiment of the invention, mutations may be detected cycle sequencing, which may include the use of a thermostable polymerase enzyme, a sequencing primer, dNTPs and limiting amounts of

-26 -

chain terminating fluorescently or radioactively labelled ddNTPs. Polyacrylamide gel electrophoresis or another technique such as capillary electrophoresis may be used to separate the products of the sequencing reactions followed by the detection of the fluorescent or radioactive labels. In one example of this embodiment of the invention, mutations in EMP2B could be determined using automated sequencing on an Applied Biosystems 3700 DNA Analyzer or 3730xl DNA Analyzer™. Mutations may be identified by comparing the sequence of a patient to that of a wildtype individual or to reference sequences found in the public databases.

10 Mutations in EMP2B may also be detected using the Invader™ genotyping system which uses a Cleavase™ Fragment Length Polymorphism (CFLP) assay as disclosed in US patent No 5,888,780 and available from Third Wave™ Technologies in Madison, Wisconsin.

Other embodiments of the invention contemplate the use of DNA chip technologies for the detection of mutations within the EMP2B gene. Among other applications, DNA chip technologies allow for the identification of mutations within the sequences of the invention through the analysis of the hybridisation patterns of a nucleic acid sample onto a high-density spatially addressable microarray of predetermined sequences. One example of a DNA chip technology suitable for the identification of mutations is the sequences of the invention is the Affymatrix GeneChip™ system, as disclosed in U.S. patent No. 6,045,996. This system uses photolithography and solid-phase chemistry to produce high density arrays containing hundreds of thousands of oligonucleotide probes. Genechips™ may be designed so as to facilitate the re-sequencing of a particular sequence, allowing for the identification of specific mutations. Computer based image analysis of the hybridisation patterns enables automatic base calling, and the incorporation of quality control measures within the analysis.

One skilled in the art will be aware that Southern blotting or Northern blotting may be used to detect pathogenic deletions or rearrangements within or near the sequences of the invention. Fluorescence *In Situ* Hybridization (FISH), fiber-FISH or other high-resolution cytogenetic methods may similarly

-27 -

be used for the detection of rearrangements or deletions that disrupt the sequences of the invention.

Another technique for the detection of mutations is denaturing HPLC analysis. Accordingly, one embodiment of the invention includes the use of a
5 Transgenomic Wave™ machine for the dHPLC analysis of nucleic acids for the identification of heterozygous mutations or polymorphisms within the sequences of the invention.

The invention also contemplates the use of mass spectroscopy for the genotyping of mutations. Mutant and wildtype nucleic acid molecules may
10 differ in mass due to the different composition of wildtype and mutant sequences, allowing for the identification of mutations on the basis of the molecular mass of different nucleotide sequences. The use of mass spectroscopy, and in particular Matrix Assisted Laser Desorption Ionisation Time of Flight (MALDI-TOF) mass spectroscopy for the genotyping of
15 mutations is well known by those skilled in the relevant art. For example, U.S. Patent No. 6,043,031 describes a fast and highly accurate mass spectrometer based process for detecting a particular nucleic acid sequence. The MassARRAY™ platform from SEQUENOM™ is an example of a commercially available system capable of genotyping single nucleotide
20 polymorphisms and detecting the mutations as described in the present invention.

(ii) Detecting the Malin Protein

In another embodiment, the present invention provides a method for detecting Lafora's disease comprising determining if the malin protein is
25 present or mutated in a sample from a mammal, preferably a human or dog, suspected of having Lafora's disease.

The malin protein of the present invention may be detected in a biological sample using antibodies that are specific for malin using various immunoassays that are discussed below. Antibodies that only react with
30 mutated malin would be useful as diagnostic agents to detect Lafora's disease. As such, antibodies would be prepared that bind only a mutated region of the protein.

-28 -

Conventional methods can be used to prepare the antibodies. For example, by using a peptide from the malin protein of the invention, polyclonal antisera or monoclonal antibodies can be made using standard methods. A mammal, (e.g., a mouse, hamster, or rabbit) can be immunized with an immunogenic form of the peptide which elicits an antibody response in the mammal. Techniques for conferring immunogenicity on a peptide include conjugation to carriers or other techniques well known in the art. For example, the peptide can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassay procedures can be used with the immunogen as antigen to assess the levels of antibodies. Following immunization, antisera can be obtained and, if desired, polyclonal antibodies isolated from the sera.

To produce monoclonal antibodies, antibody producing cells (lymphocytes) can be harvested from an immunized animal and fused with myeloma cells by standard somatic cell fusion procedures thus immortalizing these cells and yielding hybridoma cells. Such techniques are well known in the art, (e.g., the hybridoma technique originally developed by Kohler and Milstein (Nature 256, 495-497 (1975)) as well as other techniques such as the human B-cell hybridoma technique (Kozbor et al., Immunol. Today 4, 72 (1983)), the EBV hybridoma technique to produce human monoclonal antibodies (Cole et al. Monoclonal Antibodies in Cancer Therapy (1985) Allen R. Bliss, Inc., pages 77-96), and screening of combinatorial antibody libraries (Huse et al., Science 246, 1275 (1989)). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with the peptide and the monoclonal antibodies can be isolated. Therefore, the invention also contemplates hybridoma cells secreting monoclonal antibodies with specificity for a protein of the invention.

The term "antibody" as used herein is intended to include fragments thereof which also specifically react with a protein, of the invention, or peptide thereof. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above. For

-29 -

example, $F(ab')_2$ fragments can be generated by treating antibody with pepsin. The resulting $F(ab')_2$ fragment can be treated to reduce disulfide bridges to produce Fab' fragments.

Chimeric antibody derivatives, i.e., antibody molecules that combine a non human animal variable region and a human constant region are also contemplated within the scope of the invention. Chimeric antibody molecules can include, for example, the antigen binding domain from an antibody of a mouse, rat, or other species, with human constant regions. Conventional methods may be used to make chimeric antibodies containing the immunoglobulin variable region which recognizes a malin protein (see, for example, Morrison et al., Proc. Natl Acad. Sci. U.S.A. 81,6851 (1985); Takeda et al., Nature 314, 452 (1985), Cabilly et al., U.S. Patent No. 4,816,567; Boss et al., U.S. Patent No. 4,816,397; Tanaguchi et al., European Patent Publication EP171496; European Patent Publication 0173494, United Kingdom patent GB 2177096B).

Monoclonal or chimeric antibodies specifically reactive with a protein of the invention as described herein can be further humanized by producing human constant region chimeras, in which parts of the variable regions, particularly the conserved framework regions of the antigen-binding domain, are of human origin and only the hypervariable regions are of non-human origin. Such immunoglobulin molecules may be made by techniques known in the art, (e.g., Teng et al., Proc. Natl. Acad. Sci. U.S.A., 80, 7308-7312 (1983); Kozbor et al., Immunology Today, 4, 7279 (1983); Olsson et al., Meth. Enzymol., 92, 3-16 (1982)), and PCT Publication WO92/06193 or EP 0239400). Humanized antibodies can also be commercially produced (Scotgen Limited, 2 Holly Road, Twickenham, Middlesex, Great Britain.)

Specific antibodies, or antibody fragments, reactive against a protein of the invention may also be generated by screening expression libraries encoding immunoglobulin genes, or portions thereof, expressed in bacteria with peptides produced from the nucleic acid molecules of the present invention. For example, complete Fab fragments, VH regions and FV regions can be expressed in bacteria using phage expression libraries (See for

-30 -

example Ward et al., Nature 341, 544-546: (1989); Huse et al., Science 246, 1275-1281 (1989); and McCafferty et al. Nature 348, 552-554 (1990)).

Antibodies may also be prepared using DNA immunization. For example, an expression vector containing a nucleic acid of the invention (as
5 described above) may be injected into a suitable animal such as mouse. The protein of the invention will therefore be expressed *in vivo* and antibodies will be induced. The antibodies can be isolated and prepared as described above for protein immunization.

The antibodies may be labelled with a detectable marker including
10 various enzymes, fluorescent materials, luminescent materials and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, biotin, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein,
15 dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; and examples of suitable radioactive material include S-35, Cu-64, Ga-67, Zr-89, Ru-97, Tc-99m, Rh-105, Pd-109, In-111, I-123, I-125, I131, Re-186, Au-198, Au-199, Pb-203, At-211, Pb-212 and Bi-212. The antibodies may also be labelled or conjugated to one partner of a ligand
20 binding pair. Representative examples include avidin-biotin and riboflavin-riboflavin binding protein. Methods for conjugating or labelling the antibodies discussed above with the representative labels set forth above may be readily accomplished using conventional techniques.

The antibodies reactive against proteins of the invention (e.g. enzyme
25 conjugates or labelled derivatives) may be used to detect a protein of the invention in various samples, for example they may be used in any known immunoassays which rely on the binding interaction between an antigenic determinant of a protein of the invention and the antibodies. Examples of such assays are radioimmunoassays, enzyme immunoassays (e.g. ELISA),
30 immunofluorescence, immuno-precipitation, latex agglutination, hemagglutination, and histochemical tests. Thus, the antibodies may be used

-31 -

to identify or quantify the amount of a protein of the invention in a sample in order to diagnose the presence of Lafora's disease.

In a method of the invention a predetermined amount of a sample or concentrated sample is mixed with antibody or labelled antibody. The amount of antibody used in the process is dependent upon the labelling agent chosen. The resulting protein bound to antibody or labelled antibody may be isolated by conventional isolation techniques, for example, salting out, chromatography, electrophoresis, gel filtration, fractionation, absorption, polyacrylamide gel electrophoresis, agglutination, or combinations thereof.

The sample or antibody may be insolubilized, for example, the sample or antibody can be reacted using known methods with a suitable carrier. Examples of suitable carriers are Sepharose or agarose beads. When an insolubilized sample or antibody is used protein bound to antibody or unreacted antibody is isolated by washing. For example, when the sample is blotted onto a nitrocellulose membrane, the antibody bound to a protein of the invention is separated from the unreacted antibody by washing with a buffer, for example, phosphate buffered saline (PBS) with bovine serum albumin (BSA).

When labelled antibody is used, the presence of malin can be determined by measuring the amount of labelled antibody bound to a protein of the invention in the sample or of the unreacted labelled antibody. The appropriate method of measuring the labelled material is dependent upon the labelling agent.

When unlabelled antibody is used in the method of the invention, the presence of malin can be determined by measuring the amount of antibody bound to the protein using substances that interact specifically with the antibody to cause agglutination or precipitation. In particular, labelled antibody against an antibody specific for a protein of the invention, can be added to the reaction mixture. The presence of a protein of the invention can be determined by a suitable method from among the already described techniques depending on the type of labelling agent. The antibody against an antibody specific for a protein of the invention can be prepared and labelled

-32 -

- by conventional procedures known in the art which have been described herein. The antibody against an antibody specific for a protein of the invention may be a species specific anti-immunoglobulin antibody or monoclonal antibody, for example, goat anti-rabbit antibody may be used to
- 5 detect rabbit antibody specific for a protein of the invention.

(iii) Kits

- The reagents suitable for carrying out the methods of the invention may be packaged into convenient kits providing the necessary materials, packaged into suitable containers. Such kits may include all the reagents required to
- 10 detect a nucleic acid molecule or protein of the invention in a sample by means of the methods described herein, and optionally suitable supports useful in performing the methods of the invention.

- In one embodiment of the invention, the kit includes primers which are capable of amplifying a nucleic acid molecule of the invention or a
- 15 predetermined oligonucleotide fragment thereof, all the reagents required to produce the amplified nucleic acid molecule or predetermined fragment thereof in the polymerase chain reaction, and means for assaying the amplified sequences. The kit may also include restriction enzymes to digest the PCR products. In another embodiment of the invention the kit contains a
- 20 nucleotide probe which hybridizes with a nucleic acid molecule of the invention, reagents required for hybridization of the nucleotide probe with the nucleic acid molecule, and directions for its use. In a further embodiment of the invention the kit includes antibodies of the invention and reagents required for binding of the antibody to a protein of the invention in a sample.

- 25 The methods and kits of the present invention may be used to detect Lafora's disease. Samples which may be tested include bodily materials such as blood, urine, serum, tears, saliva, feces, tissues, cells and the like. In addition to human samples, samples may be taken from mammals such as non-human primates and canids such as dogs.

- 30 Before testing a sample in accordance with the methods described herein, the sample may be concentrated using techniques known in the art, such as centrifugation and filtration. For the hybridization and/or PCR-based

-33 -

methods described herein, nucleic acids may be extracted from cell extracts of the test sample using techniques known in the art.

B. *EPM2B*/Malin Modulators

In addition to antibodies and antisense oligonucleotides described
5 above, other substances that modulate *EPM2B* expression or activity may also be identified, as well as substances that modulate mutated forms of malin.

(i) Substances that Bind Malin

Substances that affect malin activity can be identified based on their
10 ability to bind to malin and/or mutated malin.

Substances which can bind with the malin of the invention may be identified by reacting the malin with a substance which potentially binds to malin, and assaying for complexes, for free substance, or for non-complexed malin, or for activation of malin. In particular, a yeast two hybrid assay system
15 may be used to identify proteins which interact with malin (Fields, S. and Song, O., 1989, Nature, 340:245-247). Systems of analysis which also may be used include ELISA.

Accordingly, the invention provides a method of identifying substances which can bind with malin, comprising the steps of:

20 (a) reacting malin and a test substance, under conditions which allow for formation of a complex between the malin and the test substance, and

(b) assaying for complexes of malin and the test substance, for free substance or for non complexed malin, wherein the presence of complexes
25 indicates that the test substance is capable of binding malin.

The malin protein used in the assay may have the amino acid sequence shown in SEQ ID NO:2 (Figure 6B) or SEQ ID NO:4 (Figure 7B) or may be a mutated protein associated with LD as described herein or may be a fragment, analog, derivative, homolog or mimetic thereof as described herein.

30 Conditions which permit the formation of substance and malin complexes may be selected having regard to factors such as the nature and amounts of the substance and the protein.

-34 -

The substance-protein complex, free substance or non-complexed proteins may be isolated by conventional isolation techniques, for example, salting out, chromatography, electrophoresis, gel filtration, fractionation, absorption, polyacrylamide gel electrophoresis, agglutination, or combinations thereof. To facilitate the assay of the components, antibody against malin or the substance, or labelled malin, or a labelled substance may be utilized. The antibodies, proteins, or substances may be labelled with a detectable substance as described above.

Malin, or the substance used in the method of the invention may be insolubilized. For example, malin or substance may be bound to a suitable carrier. Examples of suitable carriers are agarose, cellulose, dextran, Sephadex, Sepharose, carboxymethyl cellulose polystyrene, filter paper, ion-exchange resin, plastic film, plastic tube, glass beads, polyamine-methyl vinyl-ether-maleic acid copolymer, amino acid copolymer, ethylene-maleic acid copolymer, nylon, silk, etc. The carrier may be in the shape of, for example, a tube, test plate, beads, disc, sphere etc.

The insolubilized protein or substance may be prepared by reacting the material with a suitable insoluble carrier using known chemical or physical methods, for example, cyanogen bromide coupling.

The proteins or substance may also be expressed on the surface of a cell using the methods described herein.

The invention also contemplates assaying for an antagonist or agonist of the action of malin.

It will be understood that the agonists and antagonists that can be assayed using the methods of the invention may act on one or more of the binding sites on the protein or substance including agonist binding sites, competitive antagonist binding sites, non-competitive antagonist binding sites or allosteric sites.

The invention also makes it possible to screen for antagonists that inhibit the effects of an agonist of malin. Thus, the invention may be used to assay for a substance that competes for the same binding site of malin.

(ii) Peptide Mimetics

-35 -

The present invention also includes peptide mimetics of the malin and mutated malin proteins of the invention. For example, a peptide derived from a the mutated domain of malin will interact directly or indirectly with an associated molecule in such a way as to mimic the native binding of the mutated protein. Such peptides may include competitive inhibitors, enhancers, peptide mimetics, and the like. All of these peptides as well as molecules substantially homologous, complementary or otherwise functionally or structurally equivalent to these peptides may be used for purposes of the present invention.

10 "Peptide mimetics" are structures which serve as substitutes for peptides in interactions between molecules (See Morgan et al (1989), Ann. Reports Med. Chem. 24:243-252 for a review). Peptide mimetics include synthetic structures which may or may not contain amino acids and/or peptide bonds but retain the structural and functional features of a peptide, or
15 enhancer or inhibitor of the invention. Peptide mimetics also include peptoids, oligopeptoids (Simon et al (1972) Proc. Natl. Acad. Sci USA 89:9387); and peptide libraries containing peptides of a designed length representing all possible sequences of amino acids corresponding to a peptide of the invention.

20 Peptide mimetics may be designed based on information obtained by systematic replacement of L-amino acids by D-amino acids, replacement of side chains with groups having different electronic properties, and by systematic replacement of peptide bonds with amide bond replacements. Local conformational constraints can also be introduced to determine
25 conformational requirements for activity of a candidate peptide mimetic. The mimetics may include isosteric amide bonds, or D-amino acids to stabilize or promote reverse turn conformations and to help stabilize the molecule. Cyclic amino acid analogues may be used to constrain amino acid residues to particular conformational states. The mimetics can also include mimics of
30 inhibitor peptide secondary structures. These structures can model the 3-dimensional orientation of amino acid residues into the known secondary conformations of proteins. Peptoids may also be used which are oligomers of

-36 -

N-substituted amino acids and can be used as motifs for the generation of chemically diverse libraries of novel molecules.

Peptides of the invention may also be used to identify lead compounds for drug development. The structure of the peptides described herein can be readily determined by a number of methods such as NMR and X-ray crystallography. A comparison of the structures of peptides similar in sequence, but differing in the biological activities they elicit in target molecules can provide information about the structure-activity relationship of the target. Information obtained from the examination of structure-activity relationships can be used to design either modified peptides, or other small molecules or lead compounds that can be tested for predicted properties as related to the target molecule. The activity of the lead compounds can be evaluated using assays similar to those described herein.

Information about structure-activity relationships may also be obtained from co-crystallization studies. In these studies, a peptide with a desired activity is crystallized in association with a target molecule, and the X-ray structure of the complex is determined. The structure can then be compared to the structure of the target molecule in its native state, and information from such a comparison may be used to design compounds expected to possess.

20 (iii) Drug Screening Methods

In accordance with one embodiment, the invention enables a method for screening candidate compounds for their ability to increase or decrease the activity of the mutated malin protein. The method comprises providing an assay system for assaying malin activity, assaying the activity in the presence or absence of the candidate or test compound and determining whether the compound has increased or decreased malin activity. Such compounds may be useful in treating Lafora's disease.

Accordingly, the present invention provides a method for identifying a compound that affects mutated malin protein activity or expression comprising:

(a) incubating a test compound with a malin protein or a nucleic acid encoding a malin protein; and

-37 -

(b) determining an amount of malin protein activity or expression and comparing with a control (i.e. in the absence of the test substance), wherein a change in the malin protein activity or expression as compared to the control indicates that the test compound has an effect on malin protein activity or expression.

5 In accordance with a further embodiment, the invention enables a method for screening candidate compounds for their ability to increase or decrease expression of a malin protein. The method comprises putting a cell with a candidate compound, wherein the cell includes a regulatory region of a malin gene operably joined to a reporter gene coding region, and detecting a
10 change in expression of the reporter gene.

In one embodiment, the present invention enables culture systems in which cell lines which express the mutated malin gene are incubated with candidate compounds to test their effects on mutated malin expression. Such
15 culture systems can be used to identify compounds which upregulate or downregulate malin expression or its function, through the interaction with other proteins.

Such compounds can be selected from protein compounds, chemicals and various drugs that are added to the culture medium. After a period of
20 incubation in the presence of a selected test compound(s), the expression of mutated malin can be examined by quantifying the levels of malin mRNA using standard Northern blotting procedure, as described in the examples included herein, to determine any changes in expression as a result of the test compound. Cell lines transfected with constructs expressing malin can also
25 be used to test the function of compounds developed to modify the protein expression.

C. Therapeutic Uses

As previously discussed, the *EPM2B* gene and malin of the invention is likely involved in Lafora's disease. Accordingly, the present invention
30 provides a method of treating Lafora's disease comprising of administering to a cell or animal in need thereof, an effective amount of agent that modulates *EPM2B*/malin expression and/or activity. The present invention also provides

-38 -

a use of an agent that modulates *EPM2B*/malin expression and/or activity to treat Lafora's disease or to prepare a medicament to treat Lafora's disease.

The term "agent that modulates *EPM2B*/malin expression and/or activity" means any substance that can alter the expression and/or activity of the mutated *EPM2B*/malin found in the animal to be consistent with the wild type *EPM2B*/malin. Examples of agents which may be used to include administering: a nucleic acid molecule encoding wild type *EPM2B*; the wild type malin protein as well as fragments, analogs, derivatives or homologs thereof; antibodies; antisense nucleic acids; peptide mimetics; and substances isolated using the screening methods described herein that can correct the mutation to result in *EPM2B*/malin levels and/or function consistent with a person without the disease.

The term "effective amount" as used herein means an amount effective, at dosages and for periods of time necessary to achieve the desired results.

The term "animal" as used herein includes all members of the animal kingdom, including humans and dogs.

In one embodiment, the invention provides a method of treating Lafora's disease by administering to a cell or animal an effective amount of an agent that modulates the expression or the biological activity of the mutated malin protein. The present invention also provides a use of an effective amount of an agent that modulates the expression or the biological activity of the mutated malin protein to treat Lafora's disease or to prepare a medicament to treat Lafora's disease. Substances that inhibit the activity of mutated malin include peptide mimetics, malin antagonists and certain antibodies to malin. Substances that inhibit the expression of the mutated *EPM2B* gene include antisense oligonucleotides to a mutated *EPM2B* nucleic acid sequence.

In accordance with another embodiment, the present invention enables gene therapy as a potential therapeutic approach to Lafora's disease, in which normal copies of the *EPM2B* gene are introduced into patients to successfully code for normal malin protein in several different affected cell types.

-39 -

Retroviral vectors can be used for somatic cell gene therapy especially because of their high efficiency of infection and stable integration and expression. The targeted cells however must be able to divide and the expression of the levels of normal protein should be high. The full length
5 normal *EPM2B* gene can be cloned into a retroviral vector and driven from its endogenous promoter or from the retroviral long terminal repeat or from a promoter specific for the target cell type of interest. Other viral vectors which can be used include adeno-associated virus, vaccinia virus, bovine papilloma virus, or a herpesvirus such as Epstein-Barr virus. Gene transfer could also
10 be achieved using non-viral means requiring infection *in vitro*. This would include calcium phosphate, DEAE dextran, electroporation, cationic or anionic lipid formulations (liposomes) and protoplast fusion. Although these methods are available, many of these are lower efficiency.

Anti-sense based strategies can be employed to inhibit mutated
15 *EPM2B* gene function and as a basis for therapeutic drug design. The principle is based on the hypothesis that sequence specific suppression of gene expression can be achieved by intracellular hybridization between mRNA and a complementary anti-sense species. It is possible to synthesize anti-sense strand nucleotides that bind the sense strand of RNA or DNA with
20 a high degree of specificity. The formation of a hybrid RNA duplex may interfere with the processing/transport/translation and/or stability of a target mRNA.

Hybridization is required for an antisense effect to occur. Antisense effects have been described using a variety of approaches including the use
25 of antisense oligonucleotides, injection of antisense RNA, DNA and transfection of antisense RNA expression vectors.

Therapeutic antisense nucleotides can be made as oligonucleotides or expressed nucleotides. Oligonucleotides are short single strands of DNA which are usually 15 to 20 nucleic acid bases long. Expressed nucleotides
30 are made by an expression vector such as an adenoviral, retroviral or plasmid vector. The vector is administered to the cells in culture, or to a patient, whose cells then make the antisense nucleotide. Expression vectors can be

-40 -

designed to produce antisense RNA, which can vary in length from a few dozen bases to several thousand.

Antisense effects can be induced by control (sense) sequences. The extent of phenotypic changes are highly variable. Phenotypic effects induced by antisense are based on changes in criteria such as biological endpoints, protein levels, protein activation measurement and target mRNA levels.

D. Pharmaceutical Compositions

The above described substances including nucleic acids encoding *EPM2B* and mutated *EPM2B*, malin and mutated malin proteins, antibodies, and antisense oligonucleotides as well as other agents that modulate *EPM2B*/malin and/or mutated *EPM2B*/malin may be formulated into pharmaceutical compositions for administration to subjects in a biologically compatible form suitable for administration *in vivo*. By "biologically compatible form suitable for administration *in vivo*" is meant a form of the substance to be administered in which any toxic effects are outweighed by the therapeutic effects. The substances may be administered to living organisms including humans, and animals.

Administration of a therapeutically active amount of pharmaceutical compositions of the present invention is defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. For example, a therapeutically active amount of a substance may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the substance to elicit a desired response in the individual. Dosage regimes may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

An active substance may be administered in a convenient manner such as by injection (subcutaneous, intravenous, etc.), oral administration, inhalation, transdermal application, or rectal administration. Depending on the route of administration, the active substance may be coated in a material to protect the compound from the action of enzymes, acids and other natural

-41 -

conditions which may inactivate the compound. If the active substance is a nucleic acid encoding, for example, a modified *EPM2B* gene may be delivered using techniques known in the art.

The compositions described herein can be prepared by per se known methods for the preparation of pharmaceutically acceptable compositions which can be administered to subjects, such that an effective quantity of the active substance is combined in a mixture with a pharmaceutically acceptable vehicle. Suitable vehicles are described, for example, in Remington's Pharmaceutical Sciences (Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., USA 1985) or Handbook of Pharmaceutical Additives (compiled by Michael and Irene Ash, Gower Publishing Limited, Aldershot, England (1995)). On this basis, the compositions include, albeit not exclusively, solutions of the substances in association with one or more pharmaceutically acceptable vehicles or diluents, and may be contained in buffered solutions with a suitable pH and/or be iso-osmotic with physiological fluids. In this regard, reference can be made to U.S. Patent No. 5,843,456. As will also be appreciated by those skilled, administration of substances described herein may be by an inactive viral carrier.

20 E. Experimental Models

The present invention also includes methods and experimental models for studying the function of the *EPM2B* gene and malin protein. Cells, tissues and non-human animals that lack the *EPM2B* gene or partially lack in malin expression may be developed using recombinant expression vectors having a specific deletion or mutation in the *EPM2B* gene. A recombinant expression vector may be used to inactivate or alter the *EPM2B* gene by homologous recombination and thereby create an *EPM2B* deficient cell, tissue or animal.

Null alleles may be generated in cells, such as embryonic stem cells by deletion mutation. A recombinant *EPM2B* gene may also be engineered to contain an insertion mutation which inactivates *EPM2B*. Such a construct may then be introduced into a cell, such as an embryonic stem cell, by a technique such as transfection, electroporation, injection etc. Cells lacking an

-42 -

intact *EPM2B* gene may then be identified, for example by Southern blotting, Northern Blotting or by assaying for *EPM2B* using the methods described herein. Such cells may then be fused to embryonic stem cells to generate transgenic non-human animals deficient in *EPM2B*. Germline transmission of the mutation may be achieved, for example, by aggregating the embryonic stem cells with early stage embryos, such as 8 cell embryos, *in vitro*; transferring the resulting blastocysts into recipient females and; generating germline transmission of the resulting aggregation chimeras. Such a mutant animal may be used to define specific cell populations, developmental patterns and *in vivo* processes, normally dependent on *EPM2B* expression. The present invention also includes the preparation of tissue specific knock-outs of the *EPM2B* gene.

The following non-limiting example is illustrative of the present invention:

15 EXAMPLES

Example 1

The inventors examined all available databases and found 7 annotated genes in the newly defined critical region on chromosome 6p22.3 (Figure 1). Based on their predicted functional characteristics each gene was prioritized for mutation screening through DNA sequencing of LD patients carrying an *EPM2B* haplotype, but no pathogenic variants were identified. Simultaneously, the inventors' analysis led to the discovery of a previously uncharacterized apparently single-exon (1188 bp) gene sharing extensive sequence identity with orthologous units (with equivalent protein-coding potential) in other higher vertebrates (75%, 78%, 87% nucleotide identity with rat, mouse, and dog, respectively). The human gene, designated as *EPM2B*, also had at its 5'-end all of the proposed features of the consensus sequence of an eukaryotic translational initiation site and, at its 3'-end, two putative polyadenylation signals (Figure 1). Moreover, expressed sequence tag (EST) and cDNA data in human and mouse supported that the single-exon unit was a *bona fide* gene.

-43 -

EPM2B would be predicted to encode a 395 amino acid (aa) protein that the inventors have named malin (*mal* for seizure in French) containing a zinc finger of the RING type in the N-terminal half and 6 NHL-repeat domains in the C-terminal direction (Figure 1). Specifically, the zinc-binding RING-
5 finger motif (C-X₂-C-X₁₆-C-X₁-H-X₂-C-X₂-C-X₁₄-C-X₂-C) was identified (E-value of 0.0067) between residues 26-71 of malin consistent with the signature sequence (C-X₂-C-X₉₋₃₉-C-X₁₋₃-H-X₂₋₃-H-X₂-C-X₄₋₄₈-C-X₂-C) of the RING-HC type (16,17). The presence of a RING finger is predictive of an E3 ubiquitin ligase function (8,9,18). E3 ligation is the final and specific step of
10 the ubiquitin pathway transferring ubiquitin from E2, either directly or through adaptor proteins, to a specific substrate(s) to initiate its removal by the proteasome system (8). The 6 NHL domains (10-13) were predicted on the basis of presence of an approximately 44-residue motif rich in glycine and hydrophobic amino acids seeded with a cluster of charged residues (Pfam
15 detected six trusted matches for NHL domains with E-values ranging from 0.011 to 3.5) (Figure 1).

Northern-blot analysis indicated *EPM2B* is present (as at least two transcripts 1.5kb and 2.4kb in size) in all tissues examined including specific sub-regions of the brain (Figure 3). The observed transcript sizes correspond
20 near to the lengths expected between the predicted ATG-start site and the two different polyadenylation signals (Figure 1). Moreover, the expression profile was similar to that observed for *EPM2A*, both being present in all tissues in which Lafora bodies have been observed (2,19).

The complete coding region of *EPM2B* was sequenced in a cohort of
25 34 LD probands previously shown not to carry mutations in *EPM2A*. In this Example, 17 different DNA sequence alterations are described in *EPM2B* in 26 families including 8 deletions and 1 insertion leading to frame-shifts, 7 missense, and 1 non-sense change (Table 1). These mutations were found in families in both homozygous (18) and compound heterozygous (8) recessive
30 states. The four consanguineous F-C families used in the original linkage study all carried a homozygous 76T→A change producing a cysteine-to-serine alteration in one of the 7 conserved cysteine residues that are critical

-44 -

for the zinc-binding ability of the RING-finger domain (Figure 2b). The most common mutation identified (7 families) was a homozygous 205C→G transition resulting in a proline to alanine change in the RING-finger domain. The LD38324 family that was critical in refining the *EPM2B* locus (Figure 2a) carried a homozygous 2-bp deletion (1048-1049delGA) leading to a frame-shift mutation in the fifth NHL-domain (Figure 2b). Referring to Table 1, in family LD51, DNA from the proband was not available but the parents were both heterozygous carriers of 468-469delAG, which would be predicted to lead to homozygous frame-shift mutations in the LD child. All of the mutations detected would affect the putative RING or NHL motifs, or would be predicted to lead to a frame-shift or cause drastic structural change in the protein (LD483 carries a 260T→C nucleotide change which would lead to a leucine to proline alteration). Four silent DNA sequence-coding variants were identified. Three of them T312C (H104H), G372C (G124G) and T1020C (G340G) were present in five, two, and one of 100 control chromosomes, respectively. The most common polymorphism detected, C332T (P111L) (Figure 1) was observed on 42 of 100 control chromosomes.

In total, 88% of the LD families can now be accounted for by mutations in *EPM2A* (48%) and *EPM2B* (40%). The observation of 8 families with no detectable mutations in either of these genes suggests there could be additional LD loci.

Among other conditions with polyglucosan accumulation, including adult polyglucosan body disease (APBD) (20), LD is unique for the sub-cellular location of inclusions in neuronal dendrites but not axons (21), as is shown in Figure 4. The physical association of the forming polyglucosan fibrils with ER is also specific to LD (3). In APBD, which is caused by mutations in the glycogen branching enzyme (20,22,23), polyglucosans are indistinguishable in size, composition, and number from Lafora bodies, but they are located exclusively in the cell soma and axons (20). The presence of a seizure phenotype in LD but not in APBD implicates ER-associated dendritic accumulations of polyglucosans in the epilepsy of LD.

-45 -

To examine the cellular localization of malin the inventors transfected an epitope-tagged *EPM2B* construct and found that it did indeed localize at the ER and to a lesser extent within the nucleus of cultured cells (Figure 5). These results were similar to the cellular localization observed for the two alternative transcripts (A and B) of *EPM2A*, which encode isoforms of laforin found in the cytoplasm at the ER (24-26) (Figure 4b) and in the nucleus, respectively (27). The inventors most recent data implicate loss of function of the cytoplasmic form of laforin in LD based on the identification of transcript A-specific mutations (L.I. et al., manuscript submitted). Moreover, the study of murine-*EPM2A* knockouts (28) and LD patients (1,2) has shown that the ER-associated polyglucosan bodies precede or are concomitant, respectively, with onset of epilepsy.

Therefore, in the simplest explanation, LD arises due to improper clearance, and subsequent accumulation of polyglucosans in dendrites, disturbing neuronal synaptic function leading to epileptogenesis. The inventors have now shown in transgenic LD mice that laforin contacts polyglucosans (and not glycogen) providing the first physical link between a disease gene product and LD pathology (E.M.C. et al., in preparation). Laforin's only other experimentally-validated function is that of a dual-specificity phosphatase (24,25), which would predict that there is at least one phosphoprotein intermediary through which it acts. Possible candidates could be the newly discovered *EPM2AIP1* laforin-interacting protein (26), or other still to be cloned LD gene(s), or any of their interacting proteins. Malin will likely be involved via specific protein-protein interaction through its NHL domains followed by ubiquitin-mediated removal of a regulatory target(s), contributing a crucial role with laforin to safeguard neurons against Lafora bodies and epilepsy.

METHODS

Samples

All patients described in this study were formally diagnosed with adolescent-onset progressive myoclonus epilepsy based on presence of pathognomonic Lafora bodies in biopsies of skin, skeletal muscle, liver or

-46 -

brain. Each LD individual and their family members (if available) were examined for involvement of the *EPM2A* locus at 6q24 by homozygosity mapping, mutation screening, or both.

Genotyping and Mutation Screening

- 5 Information and amplification conditions on the established (*D6S274*,
D6S285, *D6S966*, *D6S1567*, *D6S1678*, *D6S1688*, *D6S1959*) and new
(*BV012563*, *BV012730*, *BV012565*, *BV012566*, *BV012568*) are found in the
UniSTS and Entrez Nucleotides database (<http://www.ncbi.nlm.nih.gov/>). DNA
Sequence variations were detected by sequencing of PCR-products. To
10 screen *EPM2B*, two sets of primer pairs that amplify overlapping fragments
were used *EPM2B*-1F: (5'-ACTGTGACCGTG ACCGAGA-3') and *EPM2B*-1R:
(CACACCCCAAGGTAAGGAGA-3'); *EPM2B*-2F: (5'-
GACTGCCATGTGGTTGTAC-3') and *EPM2B*-2R: (5'-
AAACAATTCATTAATGGCAGCA-3') (see Figure 1). PCR was performed on
15 50 ng of DNA in buffer [75 mM Tris-HCl (pH 8.8), 20 mM (NH₄)₂SO₄, 0.01%
Tween 20, 1.5 mM MgCl₂, 1M Betaine, 0.2 mM dNTP, 0.2 uM of each primer,
2.5 Units of Taq Polymerase (MBI Fermentas)]. Cycling conditions were: initial
denaturation at 94°C for 3 min followed by 35 cycles of denaturation at 94°C
for 30 sec, annealing at 60°C for 30 sec and extension at 72°C for 30 sec,
20 with a 10 min final extension at 72°C. PCR products were purified using
microCLEAN (Microzone Ltd). 3µl (100ng/µl) of purified PCR product was
used as sequencing template. For all reactions, 1µl (5 pmol) of primer, 1.5µl
5X sequencing buffer (Applied Biosystems), 1µl BigDye Terminator v3.1, and
7.5µl H₂O in a 14µl reaction volume were used. Thermocycling (MJ
25 Research, Inc.) conditions were denaturing at 96°C for 30s; annealing at 50°
for 20s; and extension at 60°C for 4 min; 35 cycles. All reactions were
subsequently purified using multiscreen-HV filter plates (Millipore) and
analyzed using an ABI-3700. All sequence variants detected in LD patients
were examined in a collection of 50 (100 chromosomes) randomly selected
30 DNA samples.

Gene identification and Northern blots

-47 -

- Gene annotation data and EST sequences were mainly obtained from the University of California Santa Cruz (UCSC) genome browser (<http://genome.ucsc.edu/>) and Celera Genomics (<http://www.celera.com/>). Additional putative genes were annotated using the Genescript algorithm
- 5 (<http://tcag.bioinfo.sickkids.on.ca/genescript/>) and multi-species VISTA (<http://www-gsd.lbl.gov/vista/>) alignments between human and mouse sequence. The RING finger domain was predicted by Pfam, Prosite, InterProScan, SMART and MotifScan. NHL domains were identified using Pfam and InterProScan. *EPM2B* orthologues were identified using BLASTN
- 10 and BLASTP analyses against the GenBank non-redundant database. Prediction of the sub-cellular localization of malin by sequence analysis was performed using PSORT II. No significant signal peptide sequence (for recognition of ER, Golgi complex, lysosome and integral plasma membrane proteins), mitochondrial targeting sequence, nuclear localization signal and
- 15 peroxisomal targeting signal was identified. The human multiple-tissue blot I and human brain blot II (Clontech) were probed with a [³²P]dCTP-labeled probe that was generated using the primers 5'-GTCACCATCACCAACGACTG-3' and 5'-TGCGAAAGACCATGAGTGAC-3', which amplified a 557bp fragment within the coding region of *EPM2B*.
- 20 Hybridization and washing conditions were performed according to the manufacturer's instructions.

Sub-cellular localization and Electron Microscopy

- The myc-tagged *EPM2A* transcript A expression construct (pcDNA3myc*EPM2A*), which encodes the cytoplasmic isoform of laforin has
- 25 been described (24). A myc-tagged *EPM2B* construct was generated (pcDNA3myc*EPM2B*) using the same general protocols. Full-length *EPM2B* was amplified by PCR from genomic DNA using the (forward) primer (5'-ggatccATGcgggccgaagc-3') containing a *Bam*HI restriction site (underlined) and the start codon (uppercase) and a (reverse) primer (5'-
- 30 gcggccgccacaattcattaatggcagac-3') containing a NotI site (underlined). This product was cloned into the corresponding sites of the mammalian expression vector pcDNA3 (Invitrogen). Myc was then introduced, in frame, after

-48 -

amplifying from a previous myc-containing vector with 5' *KpnI*-tagged and 3' *BamHI*-tagged primers. pcDNA3myc*EPM2A* and pcDNA3myc*EPM2B* (2 mg) were transfected into Cos-7 cells using Lipofectamine-Plus (Invitrogen) and exposed to lipid-DNA complex in DMEM (Sigma-Aldrich) for 5 hours. Forty-eight hours post-transfection, cultures were rinsed twice in PBS and fixed for 15 min at -20°C in an acetone:methanol (1:1) mix. They were then stained with antibodies against myc-laforin and ER marker GRP94. Cultures were blocked for 1 hour (10% BSA/PBS) and incubated with anti-Myc and anti-ER for 45 min at room temperature. Slides were washed with PBS and incubated with secondary antibody (FITC-labeled goat anti-mouse, 1:400, detectable through the green filter; Texas red-labeled donkey anti-goat, 1:400, detectable through the red filter; Jackson ImmunoResearch Laboratories) in blocking solution. Following mounting (Dako Anti-Fade), they were analyzed by immunofluorescence light microscopy. For electron microscopic examination biopsy material was obtained from the LD patient and placed into chilled Universal fixative. Using standard protocols it was then analyzed at the ultra-structural level.

Example 2

In addition to the mutations described in Example 1, the inventors have found an additional 4 mutations in *EPM2B* that is associated with Lafora's disease. These mutations were described in the same fashion as Example 1. The DNA was obtained from blood from patients with Lafora disease clinically similar to the ones described in Example 1. The three mutations in *EPM2B* are as follows: 1) A deletion of nucleotide T at position 606; 2) An A to T change at nucleotide 923; 3) A G to T change at nucleotide 580; and 4) A G to T change at nucleotide 199. The latter mutation was found in both a Sudanese and Italian family. The additional mutations are listed at the bottom of Table 1.

Example 3

Lafora's disease has also been discovered in dogs with surprising frequency. Progressive myoclonus epilepsy (PME), is common (~5%) in the popular Miniature Wirehaired Dachshund (MWHD) breed in the United

-49 -

Kingdom. The inventors characterized the clinicopathologic phenotype of these dogs and show them to have LD. The inventors mapped the underlying disease locus and identified the specific disease-associated mutation (in the *EPM2B* gene): the first coding dodecamer repeat expansion in any species and the first disease-causing tandem repeat expansion outside human. The inventors have shown that the expansion mutation is recurrent, affecting epileptic dogs other than MWHHD and demonstrate that it arises from a sequence variation particular to dogs (and other canids), predisposing the species to LD. Finally, the inventors devised a test to detect expanded alleles in carrier and presymptomatic animals and allow eradication of the disease from MWHHD and future affected canine populations.

Fourteen affected MWHHD with age of onset between six to nine years were initially evaluated. Myoclonus consisted of symmetrical split-second contractions of neck and limb muscles causing retropulsion of the dog; jaws would chatter or snap shut, and eyelids would blink rapidly. Myoclonus occurred spontaneously but could be triggered by sudden noises, or visual stimuli such as a ball rolling towards the dog or the strobe of light through trees during a car ride. Myoclonus was less likely when the dog was focused such as during chasing after rabbits and more likely with excitement or nervousness such as at feeding time or with visitors. Atonic attacks causing the animal to suddenly sit or fall and generalized seizures with rigidity, paddling, vocalization, foaming at the mouth, urination and unconsciousness were common. Older dogs were ataxic and blind, their seizures more frequent and unresponsive to medications. Eventually (ages 9 to 12), myoclonus and drop attacks were constant, compelling euthanasia. This clinical picture precisely recapitulates human LD (2), as does the underlying pathology (Figure 8).

Having confirmed that the dogs have LD, the inventors first cloned the canine *EPM2A* gene, and excluded its involvement. The inventors next undertook a genome-wide scan to localize the disease locus. The inventors genotyped 241 canine-specific microsatellite markers (31) spanning the entire canine genome in the 14 affected dogs and four unaffected relatives. The two

-50 -

markers at which the largest number of affected dogs were homozygous and the largest number of unaffected dogs heterozygous were REN94K23 and REN01G01, which map to the small 38Mb chromosome 35 (CFA35). The inventors ascertained more dogs from the extended pedigree shown in Figure 9a, and genotyped additional markers on CFA35. A maximum two-point LOD score of 2.65 at a recombination fraction $\theta=0.00$ was obtained for marker REN157J09. Multipoint linkage analysis generated a maximum LOD score of 3.38 across five markers including REN157J09 (surpassing the significance threshold of 3.2 for canine linkage studies (32) (Figure 9B), thereby linking MWHD LD to this region of CFA35.

As discussed in Example1, the inventors mapped the human *EPM2B* locus to chromosomal band 6p22 (15). Canine CFA35 is syntenic, in its entirety, to human 6p21.33-6p25.2 (31), and it was therefore possible that the same gene is mutated in human *EPM2B* patients and MWHD LD cases. The inventors cloned human *EPM2B* first (33), used its single exon to probe the RPCi-81 canine bacterial artificial chromosome (BAC) library and isolated BAC 328A12, which upon sequencing using human *EPM2B* primers revealed the complete open reading frame of canine *EPM2B* (AY560905). Radiation hybrid mapping linked *EPM2B* to marker REN157J09 (LOD score 19.3), the same marker to which MWHD LD had linked (above) with maximum LOD score (Figure 9b). In aggregate, these data argued that *EPM2B* was a viable candidate for the MWHD disease.

PCR amplification of the canine *EPM2B* gene initially failed in affected but not unaffected MWHD across a region in the 5' half of the gene. Examination of the normal sequence in this region revealed two consecutive identical dodecamers, and a third copy differing by a single nucleotide (the perfectly repeated copy is termed D and the third imperfect copy T, for ease of reference) (Figure 10a). The corresponding region in other sequenced species (Figure 10a) is not repetitive and the canine sequence is longer by 12 nucleotides, the length of one D repeat (Figure 10a). Concerted modifications of PCR conditions to amplify and sequence across this region in affected animals ultimately succeeded and revealed the MWHD LD mutation: all

-51 -

affected animals had bi-allelic expansions of the dodecamer repeat, with 19 to 26 copies of the D sequence in each allele (Figures 9a, 10b). The expansions are in-frame and might permit protein, albeit abnormal, to be made if mRNA containing them are stable.

5 The inventors next tested the effect of the expansion mutation on mRNA stability by comparing the amount of *EPM2B* mRNA in skeletal muscle from three affected dogs and two controls using quantitative RT-PCR (Supplementary Methods). *EPM2B* mRNA levels were more than 900 times reduced in the affected animals (Figure 10c).

10 The extra D sequence characterizing canine *EPM2B* was initially detected in the BAC, of Doberman Pinscher origin, and in the normal MWHD. The inventors considered whether this is a feature of only some breeds of dog, predisposing them, at least in the case of the MWHD, to expansion and L.D. The inventors sequenced *EPM2B* from two normal unrelated dogs from
15 each of 128 different breeds. Sixty percent of chromosomes had three repeats and 40% two repeats. Alleles with three repeats consisted always of two D's and one T and those with two repeats of one D and one T. Almost all breeds had examples of both variants, in homozygous or heterozygous state. Inheritance, tested in three four-generation families of different breeds, was
20 Mendelian.

Having shown that the dodecamer repeat is present across the canine species, the inventors considered whether its expansion is a cause of myoclonic seizures in other breeds of dog. The inventors tested the next non-MWHD PME case to present to the clinic, a Bassett Hound, and found a
25 homozygous seven-fold expansion (14 copies) of the D repeats (Figure 10b).

The sequence of the dodecamer expansion mutation is composed exclusively of G and C nucleotides. In the presence of the normal allele, PCR of the expanded allele was impossible, and hence carriers could not be determined. To detect carriers, the inventors deaminated the DNA, converting
30 unmethylated cytosines to uracils, and then PCR-amplified, which allowed amplification of the mutant allele, and reliable detection of carrier MWHD

-52 -

(Figure 3b). Deamination also greatly facilitated PCR amplification and diagnosis of affected animals.

The deamination method employed above was originally developed (34) for the amplification of an until now unique dodecamer repeat expansion mutation arising from a two to three-copy dodecamer repeat sequence specific to the human cystatin-B gene promoter and causing EPM1, another form of PME, by impeding cystatin-B expression (35). In humans, the more than 60 mutations in the two LD genes (*EPM2A* and *EPM2B*) (<http://projects.tcag.ca/lafora/>) are not the most common cause of PME. Instead, the single, recurring, cystatin-B promoter expansion mutation causes most human PME. To the inventors' knowledge, EPM1 has never been described in dogs, likely because they do not have the cystatin-B dodecamer repeat. LD, on the other hand is regularly reported in dogs (Basset Hounds (36,37,38), Miniature (36) and Standard (39) Poodles, Pointers (40), Corgis (41), Beagles (42,43,44) and MWH (45,46). This is likely due to recurrent expansion events of the *EPM2B* repeat, combined with the intensive inbreeding so prevalent in the domesticated dog. In other domesticated animals LD is exceedingly rare: two cows identified during mass testing for bovine spongiform encephalopathy (47) (which clinically resembles LD) and one cat (48).

Presence of the dodecamer repeat across canine breed barriers (49) suggests that its origin predates dogs and that it might therefore be present in related species. The inventors determined the distribution of the repeat in the order *Carnivora* and show that it is specific to the *Canidae* family (wolves, dogs, foxes, coyotes and jackals) (Figure 10d). The inventors were also able to discern the repeat's evolution, which occurred in two steps: appearance of the D sequence and then its duplication. Carnivores separated into *Felidae* and *Canoidae* in the Paleocene ~60 million years ago (Ma) (50,51). The D sequence is present across extant *Canoidae* but not *Felidae* (Figure 10d). It therefore first appeared after the feline-canoid split and before the subsequent divergence of canoids into *Arctoidae* and *Canidae* ~50Ma (Figure 10d). *Arctoids* flourished on the Eurasian continent where they evolved into extant

-53 -

bears, skunks, racoons, otters, etc. (50). These species have a single copy of the D sequence (Figure 10d). Canids occupied the equivalent niches on the North American landmass (successfully defending it against arctoid incursions for 40 million years) (50), and evolved by ~10Ma to present-day wolves, dogs, foxes, coyotes and jackals (51), all of which have the duplicated D sequence (Figure 10d). Duplication of the D sequence therefore occurred in the canid evolutionary line, sometime between 50Ma and 10Ma. Whether the resulting variable four-amino acid lengthening of the middle portion of malin (Figure 10a) confers a property to this E3 ligase advantageous to canids remains to be seen.

Epilepsy is the most common neurological disorder of dogs. Its prevalence is at least five times higher than in human (52). Here, the inventors have discovered the first canine epilepsy mutation. To what extent it contributes to other forms of canine epilepsy remains to be determined.

In the United Kingdom, LD is presently widespread in the MWHD, a breed with ~800 dogs registered each year. Through popular carrier sires and repeatedly bred carrier champions the disease has spread to South Africa and North America (data not shown). In the United States, different dachshund coat types are considered varieties of the same breed and are intercrossed, which might spread the disease throughout the dachshund. The mutation detection test resulting from the work described here should instead allow its eradication through controlled breeding. Meanwhile, these animals are loved pets and receive neurologic care. They far outnumber human LD patients followed in any one medical centre. This is affording the inventors important experience with best treatments for this disease, which the inventors are applying to human patients.

While the present invention has been described with reference to what are presently considered to be the preferred examples, it is to be understood that the invention is not limited to the disclosed examples. To the contrary, the invention is intended to cover various modifications and equivalent arrangements included within the spirit and scope of the appended claims.

-54 -

All publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

Table 1. *EPH2B* mutations identified in LD patients.

Family	Origin	Nucleotide change	Predicted Effect
LD6	F-Canada	Homozygous 76T→A	C26S (missense RING finger)
LD7	F-Canada	Homozygous 76T→A	C26S (missense RING finger)
LD27	F-Canada	Homozygous 76T→A	C26S (missense RING finger)
LD28	F-Canada	Homozygous 76T→A	C26S (missense RING finger)
LD23102	Brazil	Homozygous 205C→G	P69A (missense RING finger)
LDlaf100	Brazil	Homozygous 205C→G	P69A (missense RING finger)
LD5922	Italy	Homozygous 205C→G	P69A (missense RING finger)
LDlaf26	Italy	Heterozygous 205C→G 838G→A	P69A (missense RING finger) E279K (missense NHL 4)
LD41818	Spain	Heterozygous 205C→G 468delA	P69A (missense RING finger) G158fs173 (frameshift)
LD35180	F-Canada	Heterozygous 205C→G 204delC	P69A (missense RING finger) P69fs21 (frameshift RING finger)
LD29852	United States	Heterozygous 205C→G 468-469delAG	P69A (missense RING finger) G158fs16 (frameshift)
LD34477	India	Heterozygous 468-469delAG 676C→T	G158fs16 (frameshift) Q226X (nonsense NHL 3)
LD51	Brazil	Homozygous 468-469delAG	G158fs16 (frameshift)
LDlaf101	Italy	Homozygous 468-469delAG	G158fs16 (frameshift)
LDlaf9	Yugoslavia	Heterozygous 992delG 468-469delAG	G321fs2 (frameshift NHL 5) G158fs16 (frameshift)
LDlaf1	Yugoslavia	Heterozygous 992delG 468-469delAG	G321fs2 (frameshift NHL 5) G158fs16 (frameshift)
LD949	Bosnia	Homozygous 992delG	G321fs2 (frameshift NHL 5)
LD38324	Yugoslavia	Homozygous 1048-1049delGA	E340fs40 (frameshift NHL 5)
LD7635	Israel	Homozygous 373-382del10bp	T125fs103 (frameshift)
LD628	Italy	Homozygous 661-692del32bp	V16fs1 (frameshift NHL 3)
LD483	Italy	Homozygous 260T→C	L87P (missense)
LD22830	Canada	Homozygous 905A→C	Q29P (missense NHL 4)
LD32817	Pakistan	Homozygous 98T→C	F33S (missense RING finger)
LD25-9	Saudi Arabia	Homozygous 892ins2T	S298fs15 (frameshift NHL 4)
LD5487	Denmark	Heterozygous 436G→A 1100delT	D146N (missense NHL 1) V362fs20 (frameshift)
LD5489	Denmark	Homozygous 1100delT	V362fs20 (frameshift)
Laf25	Italian	606delT Homozygous	F204fs27
Laf41	Italian	923A→T Homozygous	D308V
3422	Sudanese	580G→T Homozygous	G194C
Laf44	Italian	199G→T Homozygous	E67X

FULL CITATIONS FOR REFERENCES REFERRED TO IN THE SPECIFICATION

- 5 1. Lafora, G.R., Gluck, B. Beitrag zur histopathologie der myoklonischen epilepsie. *Z Ges Neurol Psychiat* 6, 1-14 (1911).
2. Minassian, B.A. Progressive myoclonus epilepsy with polyglucosan bodies: Lafora disease. *Adv Neurol* 89, 199-210 (2002).
3. Collins, G.H., Cowden, R.R. & Nevis, A.H. Myoclonus epilepsy with Lafora
10 bodies. An ultrastructural and cytochemical study. *Arch Pathol* 86, 239-54 (1968).
4. Sakai, M., Austin, J., Witmer, F. & Trueb, L. Studies in myoclonus epilepsy (Lafora body form). II. Polyglucosans in the systemic deposits of myoclonus epilepsy and in corpora amylacea. *Neurology* 20, 160-76
15 (1970).
5. Minassian, B.A. et al. Genetic locus heterogeneity in Lafora's progressive myoclonus epilepsy. *Ann Neurol* 45, 262-5 (1999).
6. Minassian, B.A. et al. Mutations in a gene encoding a novel protein tyrosine phosphatase cause progressive myoclonus epilepsy. *Nat Genet*
20 20, 171-4 (1998).
7. Minassian, B.A. et al. Mutation spectrum and predicted function of laforin in Lafora's progressive myoclonus epilepsy. *Neurology* 55, 341-6 (2000).
8. Jackson, P.K. et al. The lore of the RINGs: substrate recognition and catalysis by ubiquitin ligases. *Trends Cell Biol* 10, 429-39 (2000).
- 25 9. Hatakeyama, S. & Nakayama, K.I. U-box proteins as a new family of ubiquitin ligases. *Biochem Biophys Res Commun* 302, 635-45 (2003).
10. Fridell, R.A., Harding, L.S., Bogerd, H.P. & Cullen, B.R. Identification of a novel human zinc finger protein that specifically interacts with the activation domain of lentiviral Tat proteins. *Virology* 209, 347-57 (1995).
- 30 11. Frank, D.J. & Roth, M.B. ncl-1 is required for the regulation of cell size and ribosomal RNA synthesis in *Caenorhabditis elegans*. *J Cell Biol* 140, 1321-9 (1998).

12. Slack, F.J. & Ruvkun, G. A novel repeat domain that is often associated with RING finger and B-box motifs. *Trends Biochem Sci* 23, 474-5 (1998).
13. Slack, F.J. et al. The lin-41 RBCC gene acts in the *C. elegans* heterochronic pathway between the let-7 regulatory RNA and the LIN-29 transcription factor. *Mol Cell* 5, 659-69 (2000).
14. Serratosa, J.M. et al. A novel protein tyrosine phosphatase gene is mutated in progressive myoclonus epilepsy of the Lafora type (EPM2A). *Hum Mol Genet* 8, 345-52 (1999).
15. Chan, E.M. et al. Genetic mapping of a new Lafora Progressive Myoclonus Epilepsy locus (EPM2B) on 6p22. *Journal of Medical Genetics*, 40, 671-5 (2003).
16. Freemont, P.S. The RING finger. A novel protein sequence motif related to the zinc finger. *Ann N Y Acad Sci* 684, 174-92 (1993).
17. Saurin, A.J., Borden, K.L., Boddy, M.N. & Freemont, P.S. Does this have a familiar RING? *Trends Biochem Sci* 21, 208-14 (1996).
18. Freemont, P.S. RING for destruction? *Curr Biol* 10, R84-7 (2000).
19. Schwarz, G.A. & Yanoff, M. Lafora disease, distinct clinico-pathological form of Unverricht's syndrome. *Arch Neurol* 12, 172-188 (1965).
20. Robitaille, Y., Carpenter, S., Karpati, G. & DiMauro, S.D. A distinct form of adult polyglucosan body disease with massive involvement of central and peripheral neuronal processes and astrocytes: a report of four cases and a review of the occurrence of polyglucosan bodies in other conditions such as Lafora's disease and normal ageing. *Brain* 103, 315-36 (1980).
21. Cavanagh, J.B. Corpora-amylacea and the family of polyglucosan diseases. *Brain Res Brain Res Rev* 29, 265-95 (1999).
22. Thon, V.J., Khalil, M. & Cannon, J.F. Isolation of human glycogen branching enzyme cDNAs by screening complementation in yeast. *J Biol Chem* 268, 7509-13 (1993).
23. Lossos, A. et al. Adult polyglucosan body disease in Ashkenazi Jewish patients carrying the Tyr329Ser mutation in the glycogen-branching enzyme gene. *Ann Neurol* 44, 867-72 (1998).

-58 -

24. Minassian, B.A. et al. Laforin is a cell membrane and endoplasmic reticulum-associated protein tyrosine phosphatase. *Ann Neurol* 49, 271-5 (2001).
25. Ganesh, S. et al. Laforin, defective in the progressive myoclonus epilepsy of Lafora type, is a dual-specificity phosphatase associated with polyribosomes. *Hum Mol Genet* 9, 2251-61 (2000).
26. Ianzano, L., Zhao, X.C., Minassian, B.A., Scherer, S.W. Identification of a novel protein interacting with laforin, the EPM2A Progressive Myoclonus Epilepsy gene product. *Genomics* 81, 579-587 (2003).
27. Ganesh, S., Suzuki, T. & Yamakawa, K. Alternative splicing modulates subcellular localization of laforin. *Biochem Biophys Res Commun* 291, 1134-7 (2002).
28. Ganesh, S. et al. Targeted disruption of the Epm2a gene causes formation of Lafora inclusion bodies, neurodegeneration, ataxia, myoclonus epilepsy and impaired behavioral response in mice. *Hum Mol Genet* 11, 1251-62 (2002).
29. Kozak, M. Interpreting cDNA sequences: some insights from studies on translation. *Mamm Genome* 7, 563-74 (1996).
30. Lovering, R. et al. Identification and preliminary characterization of a protein motif related to the zinc finger. *Proc Natl Acad Sci U S A* 90, 2112-6 (1993).
31. Guyon, R. et al. A 1-Mb resolution radiation hybrid map of the canine genome. *Proc Natl Acad Sci U S A* 100, 5296-301 (2003).
32. Gordon, D., Corwin, M. B., Mellersh, C. S., Ostrander, E. A. & Ott, J. Establishing appropriate genome-wide significance levels for canine linkage analyses. *J Hered* 94, 1-7 (2003).
33. Chan, E. M. et al. Mutations in NHLRC1 cause progressive myoclonus epilepsy. *Nat Genet* 35, 125-7 (2003).
34. Weinhaeusel, A., Morris, M. A., Antonarakis, S. E. & Haas, O. A. DNA deamination enables direct PCR amplification of the cystatin B (CSTB) gene-associated dodecamer repeat expansion in myoclonus epilepsy type Unverricht-Lundborg. *Hum Mutat* 22, 404-8 (2003).

35. Lalioti, M. D. et al. Dodecamer repeat expansion in cystatin B gene in progressive myoclonus epilepsy. *Nature* 386, 847-51 (1997).
36. Holland, J. M., Davis, W. C., Prieur, D. J. & Collins, G. H. Lafora's disease in the dog. A comparative study. *Am J Pathol* 58, 509-30 (1970).
- 5 37. Jian, Z., Alley, M. R., Cayzer, J. & Swinney, G. R. Lafora's disease in the dog. A comparative study. *N Z Vet J* 38, 75-79 (1990).
38. Kaiser, E., Krauser, K. & Schwartz-Porsche, D. [Lafora disease (progressive myoclonic epilepsy) in the Bassett hound—possibility of early diagnosis using muscle biopsy?]. *Tierarztl Prax* 19, 290-5 (1991).
- 10 39. Cusick, P. K., Cameron, A. M. & Parker, A. J. Canine neuronal glycoproteinosis-Lafora's disease in the dog. *J Am Anim Hosp Assoc* 12, 518-521 (1976).
40. Mackenzie, C. D. & Johnson, R. P. Lafora's disease in a dog. *Aust Vet J* 52, 144 (1976).
- 15 41. Davis, K. E., Finnie, J. W. & Hooper, P. T. Lafora's disease in a dog. *Aust Vet J* 67, 192-3 (1990).
42. Hegreberg, G. A. & Padgett, G. A. Inherited progressive epilepsy of the dog with comparisons to Lafora's disease of man. *Fed Proc* 35, 1202-5 (1976).
- 20 43. Montgomery, D. L. & Lee, A. C. Brain damage in the epileptic beagle dog. *Vet Pathol* 20, 160-9 (1983).
44. Gredal, H., Berendt, M. & Leifsson, P. S. Progressive myoclonus epilepsy in a beagle. *J Small Anim Pract* 44, 511-4 (2003).
45. Fitzmaurice, S. & Shelton, G. D. 'Twitchiness' in miniature wirehaired dachshunds. *Vet Rec* 143, 204 (1998).
- 25 46. Schoeman, T., Williams, J. & van Wilpe, E. Polyglucosan storage disease in a dog resembling Lafora's disease. *J Vet Intern Med* 16, 201-7 (2002).
47. Simmons, M. M. Lafora disease in the cow? *J Comp Pathol* 110, 389-401 (1994).
- 30 48. Hall, D. G., Steffens, W. L. & Lassiter, L. Lafora bodies associated with neurologic signs in a cat. *Vet Pathol* 35, 218-20 (1998).

-60 -

49. Parker, H. G. et al. Genetic structure of the purebred domestic dog.
Science 304, 1160-4 (2004).
50. Janis, C. M., Scott, K. M. & Jacobs, L. L. Evolution of Tertiary mammals of
North America (Cambridge University Press, Cambridge; New York,
5 1998).
51. Vila, C., Maldonado, J. E. & Wayne, R. K. Phylogenetic relationships,
evolution, and genetic diversity of the domestic dog. J Hered 90, 71-7
(1999).
52. Licht, B. G. et al. Clinical presentations of naturally occurring canine
10 seizures: similarities to human seizures. Epilepsy Behav 3, 460-470
(2002).

-61 -

WE CLAIM:

1. An isolated nucleic acid molecule encoding a protein with a RING-
5 finger domain and 6 NHL-motifs wherein the protein is associated with Lafora's disease.
2. A nucleic acid according to claim 1 having a sequence as shown in
SEQ ID NO:1 (Figure 6A) or SEQ ID NO:3 (Figure 7).
- 10 3. An isolated nucleic acid molecule according to claim 1 comprising
 - (a) a nucleic acid sequence as shown in SEQ ID NO:1 (Figure 6A)
or SEQ ID NO:3 (Figure 7A), wherein T can also be U;
 - (b) a nucleic acid sequence complementary to (a);
 - 15 (c) a nucleic acid sequence that has substantial sequence
homology to a nucleic acid sequence of (a) or (b);
 - (d) a nucleic acid sequence that is an analog of a nucleic acid
sequence of (a), (b) or (c); or
 - (e) a nucleic acid sequence that hybridizes to a nucleic acid
20 sequence of (a), (b), (c) or (d) under stringent hybridization conditions.
- 25 4. A method of detecting Lafora's disease comprising detecting a
mutation in a nucleic acid sequence according to any one of claims 1 to 3 in a
sample from an animal.
5. A method according to claim 4 comprising detecting a C to G change at
nucleotide number 205 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
6. A method according to claim 4 comprising detecting a T to A change at
30 nucleotide number 76 in the *EPM2B* gene sequence shown in SEQ ID NO:1.

-62 -

7. A method according to claim 4 comprising detecting a deletion of nucleotides GA at nucleotide positions 1048 and 1049 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
- 5 8. A method according to claim 4 comprising detecting a deletion of nucleotides AG at nucleotide positions 468 and 469 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
9. A method according to claim 4 comprising detecting a deletion of
10 nucleotide G at nucleotide number 992 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
10. A method according to claim 4 comprising detecting a deletion of 10 bp at nucleotide positions 373 to 382 in the *EPM2B* gene sequence shown in
15 SEQ ID NO:1.
11. A method according to claim 4 comprising detecting a deletion of 32 bp at nucleotide positions 661 to 692 in the *EPM2B* gene sequence shown in
20 SEQ ID NO:1.
12. A method according to claim 4 comprising detecting a T to C change at nucleotide number 260 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
13. A method according to claim 4 comprising detecting a A to C change at
25 nucleotide number 905 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
14. A method according to claim 4 comprising detecting a T to C change at nucleotide number 98 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
- 30 15. A method according to claim 4 comprising detecting an insert of 2 Ts at nucleotide number 892 in the *EPM2B* gene sequence shown in SEQ ID NO:1.

-63 -

16. A method according to claim 4 comprising detecting a G to A change at nucleotide number 436 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
17. A method according to claim 4 comprising detecting a deletion of
5 nucleotide T at nucleotide number 1100 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
18. A method according to claim 4 comprising detecting a deletion of
10 nucleotide T at nucleotide position 606 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
19. A method according to claim 4 comprising detecting a A to T change at nucleotide number 923 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
- 15 20. A method according to claim 4 comprising detecting a G to T change at nucleotide number 580 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
21. A method according to claim 4 comprising detecting a G to T change at nucleotide number 199 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
20
22. A method according to claim 4 comprising detecting a G to A change at nucleotide number 838 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
23. A method according to claim 4 comprising detecting a C to T change at
25 nucleotide number 676 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
24. A method according to claim 4 comprising detecting a deletion of nucleotide A at nucleotide position 468 in the *EPM2B* gene sequence shown in SEQ ID NO:1.

30

-64 -


25. A method according to claim 4 comprising detecting a deletion of nucleotide C at nucleotide position 204 in the *EPM2B* gene sequence shown in SEQ ID NO:1.
- 5 26. A method according to claim 4 comprising detecting one or more mutations in the *EPM2B* gene as indicated in Table 1.
27. A method according to claim 4 comprising detecting a repeat of the sequence GCCGCCCCCGC (SEQ ID NO:5) at nucleotide position 1001 in
10 the canine sequence of *EPM2B* shown in SEQ ID NO:3.
28. A method according to claim 22 comprising detecting at least 3 repeats of SEQ ID NO:5.
- 15 29. A method according to claim 22 comprising detecting at least 10 repeats of SEQ ID NO:5.
30. A method according to claim 22 comprising detecting from about 14 to about 26 repeats of SEQ ID NO:5.
- 20 31. A method according to any one of claims 4-26 wherein the animal is human.
32. A method according to any one of claims 4 and 26-30 wherein the
25 animal is a canid.
33. A method according to any one of claims 4 and 26-30 wherein the animal is a dog.
- 30 34. An isolated protein containing a RING-finger domain and six NHL domains which protein is associated with Lafora's disease.

-65 -

35. A protein according to claim 34 having the amino acid sequence as shown in SEQ ID NO:2 (Figure 6B) or SEQ ID NO:4 (Figure 7).
36. A method for detecting Lafora's disease comprising detecting a
5 mutation in a protein according to any one of claims 34 or 35.
37. A method according to claim 36 comprising detecting a mutation in the *EPM2B* protein as indicated in Table 1.
- 10 38. A kit for carrying out the method of any one of claims 4 to 33 or 37 comprising reagents for the detection of a mutation in a nucleic acid sequence as shown in SEQ ID NO:1 or SEQ ID NO:3.
39. A kit for carrying out the method of any one of claims 36 or 37
15 comprising reagents for the detection of a mutation in a protein sequence as shown in SEQ ID NO:2 or SEQ ID NO:5.

1/15

Fig 1



2/15

Fig 2a





Fig 2b



3/15





Fig 3

4/15

Fig 4



5/15

Fig 5a





Fig 5b



6/15

Fig 6A

1 atggcgccg aagcctcgga gagcgggcca gcgctgcatg agctcatgcg cgaggcgag
61 atcagcctgc tcgagtgcaa ggtgtgcttt gagaagtttg gccaccggca gcagcgccgc
121 ccgcgcaacc tgtcctcgcg ccacgtggtc tgcctggcct gcgtggccgc cctggcgcac
181 ccgcgcaactc tggccctcga gtgccattc tgcaggcgag ctgcccggg ctgcgacacc
241 agcgactgcc tgcgggtgct gcacctcata gagctcctgg gctcagcgct tcgcagtc
301 ccgcgccgcc atcgcgccgc cccagcgcc cccggagccc tcacctgcca ccacaccttc
361 ggcggtggtg ggaacctggt caaccccacc ggactggcgc ttgtgccaa gacggggcgt
421 gtcgtggtgg tgacgcagcg caggaggcgt gtcaagattt ttgactcagg gggaggatgc
481 gcgcatcagt ttggagagaa gggggagcgt gcccaagaca ttaggtaacc ttgtgatgtc
541 accatcacca acgactgcca tgtggtgtgc actgacgcgc gcgatcgctc catcaaatgtg
601 ttgtattttt ttggccagat caagcttgctc attggaggcc aattctcctt acctgggggt
661 gtggagacca cccctcagaa tgggattgtg gtaactgatg cggaggcagg gtccctgcac
721 ctctcggacg tcgacttcgc ggaaggggtc cttcggagaa ctgaaaggtt gcaagctcat
781 ctgtgcaatc cccgaggggt ggcagtgctc tggctcacgc gggccattgc ggtcctggag
841 caccocctg cctggggac tggggtttgc agcaccaggg tgaaagtgtt tagctcaagt
901 atgcagcttg tcggccaagt ggataccttt gggctgagcc tctactttcc ctccaaaaa
961 actgcctcgc ctgtgacctt tgatcaccag ggaatgtga ttgttcaga tacatctggt
1021 coagctatcc ttgtcttagg aaaacctgag gagtttccag taccgaagcc catggtcact
1081 catgtgtctt cgcactcgtg ggtccttacc ttcaccaagg agaattctct tctgtgtctg
1141 gacacagcat ctcatctctat aaaagtctat aaagttgact gggggtgatg ggtgtgggtg
1201 ggtccctgga atcagaagca ctagtgtgc cattaatgaa ttgtttaacc ctggataagt
1261 cacttaaaact catctatcca ggcagggata attaaaacca tctggcagac ttacaaagct
1321 tgggacagtt attggagatt aatctacat ttattgaatg catactctgt gcaaggaaat
1381 ttgcaaatat tagcttattt aatctgtact atccagttag gtaattctt ccccccaag
1441 atagagtcaa gctctgtcac ccaggctgga gtgcagaagc atgatcacag ctactacag

7/15

Fig 6A (cont'd)

1501 tttaaacgtc cccgcctcag gtggctcttc cacctoagcc toccaagtag ctgggaccac
1561 aagtgtgcat taccacactc agotaatddd tgtattttgg cagagatggg gtttcaocat
1621 gttgcccagg ctgggtctcaa actcctgagt tcaagcaatc caccttcttc ggctcccaa
1681 agtaactagga gtacaggcat agocacttgc tcagcoataa tttttattat taatctcatt
1741 gtacaagtga gaaaactgag acccagagag cttaagtac ttcctcgagg tcatagttao
1801 ttactgcott agtcccaatt tgaattcaat totgattcca aataagttgc gcttaataaa
1861 gacaacagat gtgggaaaaa tatgtgaatg tgtagtgttg ctatgtgtac tgtctttaca
1921 agtagctaat tatttttagca caaagatgtg caaagaaagg agactttatg gagagttcag
1981 gagaaaaagg atttttgtgt ggccatcaat ttoattcaat ttgcgaatgc totgatggca
2041 cattagatga agttactgtt gatcctgagt tacgtgaata agaaaaacaa ttgaactgct
2101 tattaaaaaa gtaaacatgt

8/15

Fig 6B

EPM2B protein sequence

MAAEASESGPALHELMREAEISLLECKVCFEKFQHRQRRPRNLSCGHVV
CLACVAALAHPRILALECPFCRRACRGCDTSDCLPVLHIELLGSALRQS
PAAHRAAPSAPGALTCHHTFGGWGTLVNPTGLALCPKTGRVVVVHDGRRR
VKIFDSGGGCAHQFGEKGDAAQDIRYPVDVTITNDCHVVVTDAGDRSIKV
FDFFGQIKLVIGGQFSLPWGVETTPQNGIVVTD AEGSLHLLDVDFAEGV
LRRTERLQAHLCNPRGVAVSWLTGAIAVLEHPLALGTGVCSTRVKVFSSS
MQLVGQVDTFGLSLYFPSKITASAVTFDHQGNVIVADTSGPAILCLGKPE
EFPVPKPMVTHGLSHPVALTFTKENSLVLDTASHSIKVYKVDWG

9/15

Fig 7A

Promoter (5') sequence:

```

1   CCCCAGGCC   CCCCAGGCC   CCAGGCAACC   CCAGGCCCCC   AGGCAACCCA
51  AGGCCCCCG   GCCCAAGCC   CCCCAGGTC   CCGGCCCAA   GAACCAAGCC
101 CCCCAGGCC   CCGCCCCAG   CACCAGCAC   CAAGCCCCG   CCCCCGCCG
151 CAAGCACCCA   GCCCAGCAC   CAGCCCCCG   CCCCAGCCC   AGCCCAAGCA
201 CCGAGCCCC   GCCCAGCAC   CCAGCCCCG   CACCAGCCC   CCGCCCCAGC
251 CCGAGCCCC   GTCCCCCCC   CCAGACCCA   GCCCCAGCCC   CAGCAGCAGC
301 ACCCAGCAG   GGACTGCAA   GCGTAGGCTA   CCCCAGGTGG   AACACCGTGT
351 TCTAGTTTT   CTTTCCGTT   TGCAGCTGG   GCGATCGGG   GCCACCGTCT
401 GAGCTGTGT   CCGTCCGCG   AAAGCGGAG   CGCCCCGCC   CGCCCCCGC
451 CTGCTGAAG   GTCACGGGC   TGGGCTGCG   GCGCGCGTG   CGGCCCGCGA
501 GCGTCCGCT   CGCGCCCTC   CGCAGTCAG   GCGCCGCCG   CGCGCGGGG
551 ACCGCAAGC   GCGGCCGGA   GGTGCGCGC   TGCGCCCGC   ACGTCAGGCC
601 CCGCCCCGC   CCGCCCCGC   CCGTGACCG   CCCCCGCCG   GCGCCCCGC
651 CCGCCCCGC   ACCGAGCGG   GCCGCGGGA   GCGGCGCGG   CCGCGCG

```

Coding sequence:

```

          ATG
701 GGGCCGAAG   CGCGGGGAG   CGGCGGGCG   CTGCGGGAG   TGGTCGCGA
751 GGCCGAGGT   AGCTTGCTG   AGTGCAAGT   GTGCTTCAG   AGGTTGGCC
801 ACCGCCAGC   GCGGCGCCG   CGCAACCTG   CTGCGGCCA   CGTGGTGTG
851 CTGGCTGCG   TGGCGCCCT   GGGCGACCC   CGGACGCTGG   CCCTGGAGT
901 CCCCTTCTG   GCGCGGCCCT   GCGCGGCTG   CGACACCAG   GACTGCCCTG
951 CGGTGCTTC   CCTCTGGAG   CTCTGGGCT   CGGCGCTGC   CCGAGCCCC
1001 GCGCCCCCC   GCGCGCCCC   CCGCGCGCC   CCTGCGGCC   CGGGCGCCCT
1051 CGCTGCCAT   CAGCGGTTG   GAGGCTGGG   GACCTGGTC   AACCCACGG
1101 GGCTGGCGT   GTGCCCAAG   ACCGGGCGG   TCGTGGTGT   GCACGACGG
1151 AGGAGGCGG   TCAAGATCT   TGACTCCGG   GAGGATGCG   CCATCATGT
1201 TGGAGAGAAG   GGGGAGGCT   CCCAGGACAT   TAGGTACCC   CTGAGCTCG
1251 CCGTACCAA   CGACTGCCA   GTGGTTGTA   CCGACGCGG   CGACCGTCC
1301 ATCAAAGTG   TTGATTTCT   TGCCAGATC   AAGCTCGTA   TTGGAGACA
1351 GTTTTCTTA   CTTGGGGCG   TGGAGACCA   CCCTCAGAA   GGGGTCGTG
1401 TAACTGACG   CGAGGCAAG   TCGCTGACC   TGCTGGAAG   CGACTTTGA
1451 GAAGGAGCC   TCCAGAGGAC   TGAAAAGCT   CAAGGTATC   TGTGCAACC
1501 GCGAGGGGT   GCGGTGTCT   GGCTCACTG   GGCCATTGC   GTCTGGAGC
1551 ACCCTCCGG   GCTGGGGCT   GGGGCGGCA   GCACCGCCG   GAAGGTGTC
1601 AGCCCACTA   TGCAGTGAT   CGGCCAGGT   GATACCTTG   GGCTCAGCT
1651 CTTTTTCCC   TCTAGAATA   CGGCTCCGC   CGTGACCTT   GATCACAAG
1701 GGAATGTAG   TGTTCAGAT   ACTTCTAGT   AGGCCGTCT   ATGCTTGGG
1751 CAGCTGAGG   AATTCCAGT   CTGAAGGCC   ATCATCACC   ATGGTCTTC
1801 CCATCTGTG   GCATGACCT   TCACCAAGG   GAATTCTCT   CTGTGCTGG
1851 ACAGTGACG   CCATTCCGT   AAAGTCTACA   AGGCTGACT   GGGGTAA

```

10/15

Fig 7A (cont'd)

3' UTR:

TGG
 1901 GGTGTGGTGG GGGCTCTGGA ACTGCCACTA ATCCAGTTTA ACCCTGGATG
 1951 AATTAATCCC ATCTCTCGAA CGGGGATCAT TATAACTGCC TGACAGACTT
 2001 ATAAAGGTTG AAGGTAATTA TTAAGAATA ATAATGAAGT CTACCGTTTA
 2051 TTGAGTTATG TGCTCCCTGT GCTAGGAAC TTTGCAAATA TTAGCTCAGC
 2101 GTGTCCTTAC AGTGGTACCC AGGGAGGTAA TGCCCATCAT TAATCCCATT
 2151 TTAGAGATGA GAAACTGAG ACCCGAGGGT TTAAGTGATT CTCTGAAGGT
 2201 CATGTTTACT TACTGTGACA GTCACAATGG GAACCTCTATT CTGACTCCCC
 2251 AATCCCTTGC TCCTAAGTAG GATAACAGAT GTGAGAAAA GACAGCATGT
 2301 GTCTATATGT TGTTACTGTG TGTACTCTCT TTACAGGTAG CTATTTCTCT
 2351 TGGTTGGACG TGCAGAGAAA GGAGACTTTC TAGAGAGTTC AAGAGGAAAA
 2401 AGGGTAGTGT GATGAGCATG GACGTGAGTG TCATTGAAC TGTCTGTTCT
 2451 TTGATGTCAAC AGTAGGTAGA ATGACTGTGG ATCCTTCAAC TGCCCTTGGG
 2501 AAAGGTAAAC ATGTCTGTTG GGACCTGGAT GTCCTCCATC ATAGGAAACC
 2551 AGGAAATACT AGTTGGTTGC TGCAGAAAGG CTTGTGTGGA CATAAGTTCA
 2601 AAACACTGCG CGACCACCGT ACATTCAAC ACCTCCAGTG GGAGATGGCT
 2651 GGAAGACAGT CCTGTGACAG GTCTGCATTC ATAGAACAA ANGCGCCAC
 2701 CGTTGGTTCA CGGCAGAAAT AGTTTGCTGT CCTCTTCATA ATCTGTGNCN
 2751 ACCCGAAACC CTTTGTGAT AGAGTTTTTC TCTGTGCCAT TTNAATTGT
 2801 CCCATTGCAC ACACGTGTTT CCCCTAACCA GCTCCCTTGA TGCTNAGCTA
 2851 GCATTTAGGC CACTGGTAAA CCCCTGTATA CTTCTTGAGT TGAAGTTAAG
 2901 CTTTGACCCA GATAANGNCT GCTTTAATAC NTGCAGTCGA NTGGACCGAA
 2951 TAAGGGGGAA ATTTCAGGTG AGGTGGCCGG GTTCTTTATN AACCGTTT
 3001 GGTTTGA

11/15

Fig 7B

Met Gly Ala Glu Ala Ala Gly Ser Gly Arg Ala Leu Arg Glu Leu Val
 1 5 10 15

Arg Glu Ala Glu Val Ser Leu Leu Glu Cys Lys Val Cys Phe Glu Arg
 20 25 30

Phe Gly His Arg Gln Gln Arg Arg Pro Arg Asn Leu Pro Cys Gly His
 35 40 45

Val Val Cys Leu Ala Cys Val Ala Ala Leu Ala His Pro Arg Thr Leu
 50 55 60

Ala Leu Glu Cys Pro Phe Cys Arg Arg Ala Cys Arg Gly Cys Asp Thr
 65 70 75 80

Ser Asp Cys Leu Pro Val Leu His Leu Leu Glu Leu Leu Gly Ser Ala
 85 90 95

Leu Arg Pro Ala Pro Ala Ala Pro Arg Ala Ala Pro Arg Ala Ala Pro
 100 105 110

Cys Ala Pro Gly Ala Leu Ala Cys His His Ala Phe Gly Gly Trp Gly
 115 120 125

Thr Leu Val Asn Pro Thr Gly Leu Ala Leu Cys Pro Lys Thr Gly Arg
 130 135 140

Val Val Val Val His Asp Gly Arg Arg Arg Val Lys Ile Phe Asp Ser
 145 150 155 160

Gly Gly Gly Cys Ala His Gln Phe Gly Glu Lys Gly Glu Ala Ala Gln
 165 170 175

Asp Ile Arg Tyr Pro Leu Asp Val Ala Val Thr Asn Asp Cys His Val
 180 185 190

Val Val Thr Asp Ala Gly Asp Arg Ser Ile Lys Val Phe Asp Phe Phe
 195 200 205

Gly Gln Ile Lys Leu Val Ile Gly Asp Gln Phe Ser Leu Pro Trp Gly
 210 215 220

12/15

Fig 7B (cont'd)

Val Glu Thr Thr Pro Gln Asn Gly Val Val Val Thr Asp Ala Glu Ala
 225 230 235 240

Gly Ser Leu His Leu Leu Glu Val Asp Phe Ala Glu Gly Ala Leu Gln
 245 250 255

Arg Thr Glu Lys Leu Gln Gly His Leu Cys Asn Pro Arg Gly Val Ala
 260 265 270

Val Ser Trp Leu Thr Gly Ala Ile Ala Val Leu Glu His Pro Pro Gly
 275 280 285

Leu Gly Ala Gly Ala Gly Ser Thr Ala Val Lys Val Phe Ser Pro Thr
 290 295 300

Met Gln Leu Ile Gly Gln Val Asp Thr Phe Gly Leu Ser Leu Phe Phe
 305 310 315 320

Pro Ser Arg Ile Thr Ala Ser Ala Val Thr Phe Asp His Gln Gly Asn
 325 330 335

Val Ile Val Ala Asp Thr Ser Ser Gln Ala Val Leu Cys Leu Gly Gln
 340 345 350

Pro Glu Glu Phe Pro Val Leu Lys Pro Ile Ile Thr His Gly Leu Ser
 355 360 365

His Pro Val Ala Leu Thr Phe Thr Lys Glu Asn Ser Leu Leu Val Leu
 370 375 380

Asp Ser Ala Ala His Ser Val Lys Val Tyr Lys Ala Asp Trp Gly
 385 390 395

13/15




Fig 8

1/13

SEQUENCE LISTING

<110> The Hospital for Sick Children

Scherer, Stephen W.

Minassian, Berge A.

<120> Novel Lafora's Disease Gene

<130> 9962-58

<150> US 60/491,968

<151> 2003-08-04

<160> 5

<170> PatentIn version 3.1

<210> 1

<211> 2120

<212> DNA

<213> Homo sapiens

```

<400> 1
atggcggcgg aagcctcggg gagcggggcca gcgctgcatt agctcatgcg cgaggcggag      60
atcagcctgc tcgagtgcaa ggtgtgcttt gagaagtatt gccaccggca gcagcggcgc      120
ccgcgcaacc tgctctcgcg ccactgggtc tgctgggctt gcgtggccgc cctggcgcac      180
ccgcgcactc tggccctcga gtgcccatc tcgaggcgag ctgtccgggg ctgcgacacc      240
agcgactgcg tgcgggtgct gcacctcata gagctcctgg gctcagcgct tcgccagctc      300
ccggccgccc atcgcgccgc cccacggccc ccgggagccc tcacctgcca ccacaccttc      360
ggcggctggg ggaccctggt caacccacc ccgactggcg tttgtcccaa gacggggcgt      420

```

2/13

gtcgtggtgg tgcacgacgg caggaggcgt gtcaagattt ttgactcagg gggaggatgc	480
gcgcatcagt ttggagagaa gggggacgct gcccaagaca ttaggtaccc tgtggtgtc	540
accatcacca acgactgcc tgtggtgtc actgacgccc gcgactcgctc catcaaagt	600
tttgattttt ttggccagat caagcttgct attggaggcc aattctcctt accttgggggt	660
gtggagacca cccctcagaa tgggattgtg gtaactgatg cggaggcagg gtccctgcac	720
ctcctggagc tcgacttcgc ggaaggggtc cttcggagaa ctgaaaggtt gcaagctcat	780
ctgtgacaat ccgaggggtt ggcagtgctt tggctcacgg gggccattgc ggtcctggag	840
caccccctgg ccctggggac tggggtttgc agcaccaggg tgaaagtgtt tagctcaagt	900
atgcagcttg tcggccaagt ggataccttt gggctgagcc tctactttcc tccaaaata	960
actgcctcgg ctgtgacctt tgatcaccag ggaagtgtga ttgttgacga tacactcgtt	1020
ccagctatcc ttgtcttagg aaaacctgag gagtttcagg taccgaagcc catggtcact	1080
catggtcttt cgcactctgt ggctcttacc ttcaccaagg agaattctct tcttgtgtg	1140
gacacgcat ctcatctcat aaaagtctat aaagttagct gggggtgatg ggtcgggtg	1200
ggtccctgga atcagaagca ctagtgtgc cattaatgaa ttgtttaacc ctggataagt	1260
cacttaact catctatcca ggcagggata attaaaacca tctggcagac ttacaaaagt	1320
tgggcagctt attggagatt aatctaccat ttattgaatg catactctgt gcaaggaaat	1380
ttgcaaatat tagcttattt aatctgtact atccagttag gtaatttctt ccccccaag	1440
atagagtcaa gctctgtcac ccaggctgga gtgcagaagc atgacacag ctactacag	1500
tttcaacgtc ccccgctcag gtggtccttc cactcagcc tcccaagtag ctgggaccac	1560
aagtgtgcat taccacactc agctaatttt tgtattttgg cagagatggg gtttcccat	1620
gttccccagg ctggtctcaa actcctgagt tcaagcaatc cacttcctc ggcctcccaa	1680
agtactagga gtacaggcat agccacttgc tcagccataa tttttattat taatctcatt	1740
gtacaagtga gaaaactgag acccagagag cttaaagtac ttccctcgagg tcatagttac	1800
ttactgcctt agtcccaatt tgaattcaat tctgattcca aataagttgc gcttaataa	1860
gacaacagat gtgggaaaaa tatgtgaatg tgtagtgttg ctatgtgtac tgtctttaca	1920
agtagcta at ttttttagca caaagatgtg caaagaaagg agactttatg gagagtacg	1980
gagaaaaagg atttttgtgtt ggccatcact ttcattcaat ttgcgactgc tctgatggca	2040
cattagatga agttactgtt gatcctgagt tacgtgaata agaaaaacaa ttgaactgct	2100
tattaaaaaa gtaaacatgt	2120

3/13

<210> 2

<211> 395

<212> PRT

<213> Homo sapiens

<400> 2

Met Ala Ala Glu Ala Ser Glu Ser Gly Pro Ala Leu His Glu Leu Met
 1 5 10 15

Arg Glu Ala Glu Ile Ser Leu Leu Glu Cys Lys Val Cys Phe Glu Lys
 20 25 30

Phe Gly His Arg Gln Gln Arg Arg Pro Arg Asn Leu Ser Cys Gly His
 35 40 45

Val Val Cys Leu Ala Cys Val Ala Ala Leu Ala His Pro Arg Thr Leu
 50 55 60

Ala Leu Glu Cys Pro Phe Cys Arg Arg Ala Cys Arg Gly Cys Asp Thr
 65 70 75 80

Ser Asp Cys Leu Pro Val Leu His Leu Ile Glu Leu Leu Gly Ser Ala
 85 90 95

Leu Arg Gln Ser Pro Ala Ala His Arg Ala Ala Pro Ser Ala Pro Gly
 100 105 110

Ala Leu Thr Cys His His Thr Phe Gly Gly Trp Gly Thr Leu Val Asn
 115 120 125

Pro Thr Gly Leu Ala Leu Cys Pro Lys Thr Gly Arg Val Val Val Val
 130 135 140

His Asp Gly Arg Arg Arg Val Lys Ile Phe Asp Ser Gly Gly Gly Cys
 145 150 155 160

Ala His Gln Phe Gly Glu Lys Gly Asp Ala Ala Gln Asp Ile Arg Tyr
 165 170 175

Pro Val Asp Val Thr Ile Thr Asn Asp Cys His Val Val Val Thr Asp
 180 185 190

4/13

Ala Gly Asp Arg Ser Ile Lys Val Phe Asp Phe Phe Gly Gln Ile Lys
195 200 205

Leu Val Ile Gly Gly Gln Phe Ser Leu Pro Trp Gly Val Glu Thr Thr
210 215 220

Pro Gln Asn Gly Ile Val Val Thr Asp Ala Glu Ala Gly Ser Leu His
225 230 235 240

Leu Leu Asp Val Asp Phe Ala Glu Gly Val Leu Arg Arg Thr Glu Arg
245 250 255

Leu Gln Ala His Leu Cys Asn Pro Arg Gly Val Ala Val Ser Trp Leu
260 265 270

Thr Gly Ala Ile Ala Val Leu Glu His Pro Leu Ala Leu Gly Thr Gly
275 280 285

Val Cys Ser Thr Arg Val Lys Val Phe Ser Ser Ser Met Gln Leu Val
290 295 300

Gly Gln Val Asp Thr Phe Gly Leu Ser Leu Tyr Phe Pro Ser Lys Ile
305 310 315 320

Thr Ala Ser Ala Val Thr Phe Asp His Gln Gly Asn Val Ile Val Ala
325 330 335

Asp Thr Ser Gly Pro Ala Ile Leu Cys Leu Gly Lys Pro Glu Glu Phe
340 345 350

Pro Val Pro Lys Pro Met Val Thr His Gly Leu Ser His Pro Val Ala
355 360 365

Leu Thr Phe Thr Lys Glu Asn Ser Leu Leu Val Leu Asp Thr Ala Ser
370 375 380

His Ser Ile Lys Val Tyr Lys Val Asp Trp Gly
385 390 395

<210> 3

<211> 3008

<212> DNA

5/13

<213> Canis sp.

<220>

<221> CDS

<222> (698)..(1897)

<223>

<220>

<221> misc_feature

<222> (2692)..(2692)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2748)..(2748)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2750)..(2750)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2793)..(2793)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2845)..(2845)

6/13

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2916)..(2916)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2918)..(2918)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2931)..(2931)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2941)..(2941)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2990)..(2990)

<223> N=any nucleic acid

<400> 3

ccccaaaggcc ccccgggccc ccaggcaacc ccaggccccc aggcaaccca agggccccccy 60

gccccaaaggcc ccccgagttc ccggcccacaa gaaccaaggcc ccccgggccc ccggccccag 120

7/13

cacccagcac caagcccccg ccccccgccc caagcaccga gcccagcac ccagcccccg	180
ccccagcccc agccccagca cccagccccc gcccagcac ccagccccag caccagcccc	240
ccgccccagc cccagccccc gtcccccccc ccagcaccga gcccagcccc cagcagcagc	300
accagcaggg ggactgcaaa gcgtaggcta cccaggtgg aacacgtgt tctagttttg	360
ctttgcggtt tgcagcctgg gcgatcgggg gccaccgctc gagcctgttt cccgtcgcgg	420
aaagcggagc cgccccgcc cgccccccc ctgectgaag gtcaccggcc tgggctgtcg	480
gcgcgcgggtg cggccccga gcgtccgctc ccgcgccctc cgcagtcagc gcccgccgc	540
ccgcgggggg accgcaggcc gcggccgaga ggctgcgcgc tgcgccgcgc acgtcaggcc	600
ccgccccgcc ccgccccgcc ccgtgacggg ccccgggccc ggccccggcc ccggccccgg	660
accgagcggc gcccgggga gcggcgggcg ccgcgcg atg ggg gcc gaa gcg gcg	715
Met Gly Ala Glu Ala Ala	
1 5	
ggg agc ggg cgg gcg ctg cgg gag ctg gtg cgc gag gcc gag gtc agc	763
Gly Ser Gly Arg Ala Leu Arg Glu Leu Val Arg Glu Ala Glu Val Ser	
10 15 20	
ttg ctc gag tgc aag gtg tgc ttc gag agg ttc ggc cac cgc cag cag	811
Leu Leu Glu Cys Lys Val Cys Phe Glu Arg Phe Gly His Arg Gln Gln	
25 30 35	
cgg cgc ccg cgc aac ctg ccc tgc ggc cac gtg. gtg tgc ctg gcc tgc	859
Arg Arg Pro Arg Asn Leu Pro Cys Gly His Val Val Cys Leu Ala Cys	
40 45 50	
gtg gcg gcc ctg gcg cac ccg cgg acg ctg gcc ctg gag tgc ccc ttc	907
Val Ala Ala Leu Ala His Pro Arg Thr Leu Ala Leu Glu Cys Pro Phe	
55 60 65 70	
tgc cgc cgg gcc tgc cgc ggc tgc gac acc agc gac tgc ctg ccg gtg	955
Cys Arg Arg Ala Cys Arg Gly Cys Asp Thr Ser Asp Cys Leu Pro Val	
75 80 85	
ctt cac ctc ctg gag ctc ctg ggc tgc gcg ctg cgc cca gcc ccc gcc	1003
Leu His Leu Leu Glu Leu Leu Gly Ser Ala Leu Arg Pro Ala Pro Ala	
90 95 100	
gcc ccc cgc gcc gcc ccc cgc gcc gcc ccc tgc gcc ccg ggc gcc ctc	1051
Ala Pro Arg Ala Ala Pro Arg Ala Ala Pro Cys Ala Pro Gly Ala Leu	
105 110 115	
gcc tgc cat cac gcg ttc gga ggc tgg ggg acc ctg gtc aac ccc acg	1099
Ala Cys His His Ala Phe Gly Gly Trp Gly Thr Leu Val Asn Pro Thr	
120 125 130	
ggg ctg gcg ctg tgc ccc aag acc ggg cgg gtc gtg gtg gtg cac gac	1147
Gly Leu Ala Leu Cys Pro Lys Thr Gly Arg Val Val Val Val His Asp	
135 140 145 150	

8/13

ggc agg agg cgg gtc aag atc ttt gac tcc ggg gga gga tgc gcc cat Gly Arg Arg Arg Val Lys Ile Phe Asp Ser Gly Gly Gly Cys Ala His 155 160 165	1195
cag ttt gga gag aag ggg gag gct gcc cag gac att agg tac ccc ctg Gln Phe Gly Glu Lys Gly Glu Ala Ala Gln Asp Ile Arg Tyr Pro Leu 170 175 180	1243
gac gtc gcc gtc acc aac gac tgc cac gtg gtt gtc acc gac gcc ggc Asp Val Ala Val Thr Asn Asp Cys His Val Val Val Thr Asp Ala Gly 185 190 195	1291
gac cgc tcc atc aaa gtg ttt gat ttc ttt ggc cag atc aag ctc gtc Asp Arg Ser Ile Lys Val Phe Asp Phe Phe Gln Ile Lys Leu Val 200 205 210	1339
att gga gac cag ttt tcc tta cct tgg ggc gtg gag acc acc cct cag Ile Gly Asp Gln Phe Ser Leu Pro Trp Gly Val Glu Thr Thr Pro Gln 215 220 225 230	1387
aat ggg gtc gtg gta act gac gcc gag gca ggg tgc ctg cac ctg ctg Asn Gly Val Val Val Thr Asp Ala Glu Ala Gly Ser Leu His Leu Leu 235 240 245	1435
gaa gtc gac ttt gca gaa gga gcc ctc cag agg act gaa aag ctg caa Glu Val Asp Phe Ala Glu Gly Ala Leu Gln Arg Thr Glu Lys Leu Gln 250 255 260	1483
ggt cat ctg tgc aac ccg cga ggg gtg gcc gtg tcc tgg etc act ggg Gly His Leu Cys Asn Pro Arg Gly Val Ala Val Ser Trp Leu Thr Gly 265 270 275	1531
gcc att gcg gtc ctg gag cac cct ccg ggg ctg ggg gct ggg gcg ggc Ala Ile Ala Val Leu Glu His Pro Pro Gly Leu Gly Ala Gly Ala Gly 280 285 290	1579
agc acc gcc gtg aag gtg ttc agc cca act atg cag ctg atc gcc cag Ser Thr Ala Val Lys Val Phe Ser Pro Thr Met Gln Leu Ile Gly Gln 295 300 305 310	1627
gtg gat acc ttt ggg ctc agc ctc ttt ttc ccc tct aga ata acc gcc Val Asp Thr Phe Gly Leu Ser Leu Phe Phe Pro Ser Arg Ile Thr Ala 315 320 325	1675
tcc gcc gtg acc ttt gat cac cag ggg aat gtg att gtt gca gat act Ser Ala Val Thr Phe Asp His Gln Gly Asn Val Ile Val Ala Asp Thr 330 335 340	1723
tct agt cag gcc gtc cta tgc ttg gga cag cct gag gaa ttt cca gtc Ser Ser Gln Ala Val Leu Cys Leu Gly Gln Pro Glu Glu Phe Pro Val 345 350 355	1771
ctg aag ccc atc atc acc cat ggt ctt tcc cat cct gtg gca ctg acc Leu Lys Pro Ile Ile Thr His Gly Leu Ser His Pro Val Ala Leu Thr 360 365 370	1819
ttc acc aag gag aat tct ctt ctt gtg ctg gac agt gca gcc cat tcc Phe Thr Lys Glu Asn Ser Leu Leu Val Leu Asp Ser Ala Ala His Ser 375 380 385 390	1867

9/13

gta aaa gtc tac aag gct gac tgg ggg taa tgggggtgtgg tgggggtcct 1917
Val Lys Val Tyr Lys Ala Asp Trp Gly
395
ggaactgccca ctaatccagt ttaaccctgg atgaattaat cccatctctc gaacggggat 1977
cattataact gcctgcagaca cttataaagg ttgaaggtaa ttattaaaga ataataatga 2037
agtctaccgt ttattgagtt atgtgctccc tgtgctagga aactttgcaa atattagctc 2097
agcgtgtcct tacagtggtta ccaggggagg taatgcccat cattaatccc attttagaga 2157
tgagaaaact gagaccgag ggtttaagtg attctctgaa ggtcatgttt acttactgtg 2217
acagtcacaa tgggaactct attctgactc cccaatccct tgcctcctag taggataaca 2277
gatgtgagaa aacgcacaga tgtgtctata tgttgttact gtgtgtactc tctttacagg 2337
tagctatttc tcttggttgg acgtgcagag aaaggagact ttctagagag tccaagagga 2397
aaaagggtag tgtgatgagc atggacgtga gtgtcattga acttgctggg tctttgatgt 2457
cacagtaggt agaatgactg tggatccttc aactgccctt gggaaaggta aacatgtctg 2517
ttgggacctg gatgtcctcc atcataggaa cccaggaaat actagtgtgt tgcgtgcaga 2577
aggcttgtgt ggacataagt tcaaaactac tgcgcaccac cgtacattca cacacctcca 2637
gtgggagatg gctggaagac agtcctgtga caggctgtca ttcatagaac aagangccgc 2697
cacogttggt tcacggcaga atgagtttgc ctgcctcttc ataacttgtg ncnaccgcga 2757
acccttttgt gatagagttt ttctctgtgc cattttnaatt tgtccattg cacacactgt 2817
tttccctaa ccagctccct tgatgctnag cttagcattta ggccactggt aaaccctgt 2877
atactctctg agttgaagtt aagctttgac ccagataang nctgctttaa taentgcagt 2937
cgantggacc gaataagggg gaaatttcag gtgaggtggc cgggttcttt atnaaccggt 2997
tttggtttgt a 3008

<210> 4

<211> 399

<212> PRT

<213> Canis sp.

<220>

<221> misc_feature

<222> (2692)..(2692)

10/13

<223> N=any nucleic acid
<220>
<221> misc_feature
<222> (2748)..(2748)
<223> N=any nucleic acid
<220>
<221> misc_feature
<222> (2750)..(2750)
<223> N=any nucleic acid
<220>
<221> misc_feature
<222> (2793)..(2793)
<223> N=any nucleic acid
<220>
<221> misc_feature
<222> (2845)..(2845)
<223> N=any nucleic acid
<220>
<221> misc_feature
<222> (2916)..(2916)
<223> N=any nucleic acid
<220>
<221> misc_feature
<222> (2918)..(2918)
<223> N=any nucleic acid
<220>
<221> misc_feature
<222> (2931)..(2931)
<223> N=any nucleic acid

11/13

<220>

<221> misc_feature

<222> (2941)..(2941)

<223> N=any nucleic acid

<220>

<221> misc_feature

<222> (2990)..(2990)

<223> N=any nucleic acid

<400> 4

Met Gly Ala Glu Ala Ala Gly Ser Gly Arg Ala Leu Arg Glu Leu Val
 1 5 10 15

Arg Glu Ala Glu Val Ser Leu Leu Glu Cys Lys Val Cys Phe Glu Arg
 20 25 30

Phe Gly His Arg Gln Gln Arg Arg Pro Arg Asn Leu Pro Cys Gly His
 35 40 45

Val Val Cys Leu Ala Cys Val Ala Ala Leu Ala His Pro Arg Thr Leu
 50 55 60

Ala Leu Glu Cys Pro Phe Cys Arg Arg Ala Cys Arg Gly Cys Asp Thr
 65 70 75 80

Ser Asp Cys Leu Pro Val Leu His Leu Leu Glu Leu Leu Gly Ser Ala
 85 90 95

Leu Arg Pro Ala Pro Ala Ala Pro Arg Ala Ala Pro Arg Ala Ala Pro
 100 105 110

Cys Ala Pro Gly Ala Leu Ala Cys His His Ala Phe Gly Gly Trp Gly
 115 120 125

Thr Leu Val Asn Pro Thr Gly Leu Ala Leu Cys Pro Lys Thr Gly Arg
 130 135 140

Val Val Val Val His Asp Gly Arg Arg Arg Val Lys Ile Phe Asp Ser
 145 150 155 160

Gly Gly Gly Cys Ala His Gln Phe Gly Glu Lys Gly Glu Ala Ala Gln

12/13

165

170

175

Asp Ile Arg Tyr Pro Leu Asp Val Ala Val Thr Asn Asp Cys His Val
 180 185 190

Val Val Thr Asp Ala Gly Asp Arg Ser Ile Lys Val Phe Asp Phe Phe
 195 200 205

Gly Gln Ile Lys Leu Val Ile Gly Asp Gln Phe Ser Leu Pro Trp Gly
 210 215 220

Val Glu Thr Thr Pro Gln Asn Gly Val Val Val Thr Asp Ala Glu Ala
 225 230 235 240

Gly Ser Leu His Leu Leu Glu Val Asp Phe Ala Glu Gly Ala Leu Gln
 245 250 255

Arg Thr Glu Lys Leu Gln Gly His Leu Cys Asn Pro Arg Gly Val Ala
 260 265 270

Val Ser Trp Leu Thr Gly Ala Ile Ala Val Leu Glu His Pro Pro Gly
 275 280 285

Leu Gly Ala Gly Ala Gly Ser Thr Ala Val Lys Val Phe Ser Pro Thr
 290 295 300

Met Gln Leu Ile Gly Gln Val Asp Thr Phe Gly Leu Ser Leu Phe Phe
 305 310 315 320

Pro Ser Arg Ile Thr Ala Ser Ala Val Thr Phe Asp His Gln Gly Asn
 325 330 335

Val Ile Val Ala Asp Thr Ser Ser Gln Ala Val Leu Cys Leu Gly Gln
 340 345 350

Pro Glu Glu Phe Pro Val Leu Lys Pro Ile Ile Thr His Gly Leu Ser
 355 360 365

His Pro Val Ala Leu Thr Phe Thr Lys Glu Asn Ser Leu Leu Val Leu
 370 375 380

Asp Ser Ala Ala His Ser Val Lys Val Tyr Lys Ala Asp Trp Gly
 385 390 395

13/13

<210> 5

<211> 12

<212> DNA

<213> Canis sp.

<400> 5

gccgcccccc gc

12

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2004/001449

A. CLASSIFICATION OF SUBJECT MATTER

C12N 15/12 C07K 14/47 C12Q 1/68 G01N 33/88

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N 15/12 C07K 14/47 C12Q 1/68

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base, and, where practicable, search terms used)

GenBank (incl. EST and HTG), PIR, UNIPROT, Dgene, Delphion, CaPlus

Keywords: EPM2B, Lafora, malin Sequence ID numbers 1 to 4

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	CHEN, E. M. et al., "Mutations in NHLRC1 cause progressive myoclonus epilepsy", NATURE GENETICS. October 2003 (Epub 07 September 2003), Vol. 35, No. 2, pages 125-127. whole document.	1 to 39
X Y	BLAKEY, S. "Human DNA sequence from clone RP11-204B7 on chromosome 6, complete sequence". GENBANK database. 27 April 2001. Accession number AL589723. DNA sequence nucleic acids 85181 to 83042 (minus strand). DNA sequence nucleic acids 84858 to 83960 (minus strand).	1 to 3 34 and 35
X	"Mus musculus adult male corpora quadrigemina cDNA, RIKEN full-length enriched library, clone, B230309E09 product: hypothetical NHL repeat/RING finger containing protein, full insert sequence". GENBANK database 05 December 2002. Accession number AK045746 Protein sequence amino acids 16 to 395. Protein sequence amino acids 1 to 394.	1, 3 and 34
Y, P	"Hypothetical protein CBG06802 [Caenorhabditis briggsae]". GENBANK database 24 November 2003. Accession number CAE62684. Whole protein sequence	1 and 34
A, P	CHAN, E. M. et al., "Genetic mapping of a new Lafora progressing myoclonus	1 to 39

Further documents are listed in the continuation of Box C.

Patent family members are listed in annex.

- * Special categories of cited documents: "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "A" document defining the general state of the art which is not considered to be of particular relevance "X"
- "E" earlier application or patent but published on or after the international filing date "Y"
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "Y"
- "O" document referring to an oral disclosure, use, exhibition or other means "A"
- "P" document published prior to the international filing date but later than the priority date claimed

document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

document member of the same patent family

Date of the actual completion of the international-type search
19 October 2004Date of mailing of the international-type search report
14 January 2005 (14-01-2005)

Name and mailing address of the ISA/
Commissioner of Patents
Canadian Patent Office - PCT
Ottawa/Gatineau K1A 0C9
Facsimile No. 1-819-953-9358

Authorized officer
Kathleen Pound (819) 953-9757

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2004/001449

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MINASSIAN, B. A. et al., "Mutations in a gene encoding a novel protein tyrosine phosphatase cause progressive myoclonus epilepsy". NATURE GENETICS. October 1998, Vol. 20, No. 2, pages 171-174.	1 to 39
A	LEHESIÖKI, A.-E. "Molecular background of progressive myoclonus epilepsy" EMBO J. July 2003 Vol. 22, No. 14, pages 3473-3478.	1 to 39